

On Parameter Estimation and Hypothesis Testing for Poisson Processes in Case of a Change-point with Variable Jump Size

(joint work with Lin YANG)

Sergueï DACHIAN
Université de Lille 1
Lille, France



Model of observations



Model of observations

Recall: $X = (X(t), t \geq 0)$ is an (inhomogeneous) **Poisson process** with **intensity function** $\lambda(t), t \geq 0$, if $X_0 = 0$ and the increments of X on disjoint intervals are independent Poisson random variables:

$$\mathbf{P}\{X(t) - X(s) = k\} = \frac{\left(\int_s^t \lambda(t) dt\right)^k}{k!} \exp \left\{ - \int_s^t \lambda(t) dt \right\}.$$



Model of observations

Recall: $X = (X(t), t \geq 0)$ is an (inhomogeneous) **Poisson process** with **intensity function** $\lambda(t), t \geq 0$, if $X_0 = 0$ and the increments of X on disjoint intervals are independent Poisson random variables:

$$\mathbf{P}\{X(t) - X(s) = k\} = \frac{\left(\int_s^t \lambda(t) dt\right)^k}{k!} \exp \left\{ - \int_s^t \lambda(t) dt \right\}.$$

Observations: n independent realizations $X^{(n)} = (X_1^{(n)}, \dots, X_n^{(n)})$ of a Poisson process with intensity function $\lambda_{\vartheta}^{(n)}(t), 0 \leq t \leq \tau$.



Model of observations

Recall: $X = (X(t), t \geq 0)$ is an (inhomogeneous) **Poisson process** with **intensity function** $\lambda(t), t \geq 0$, if $X_0 = 0$ and the increments of X on disjoint intervals are independent Poisson random variables:

$$\mathbf{P}\{X(t) - X(s) = k\} = \frac{\left(\int_s^t \lambda(t) dt\right)^k}{k!} \exp \left\{ - \int_s^t \lambda(t) dt \right\}.$$

Observations: n independent realizations $X^{(n)} = (X_1^{(n)}, \dots, X_n^{(n)})$ of a Poisson process with intensity function $\lambda_{\vartheta}^{(n)}(t), 0 \leq t \leq \tau$.

Unknown parameter: $\vartheta \in \Theta = (\alpha, \beta), 0 \leq \alpha < \beta \leq \tau$.



Model of observations

Recall: $X = (X(t), t \geq 0)$ is an (inhomogeneous) **Poisson process** with **intensity function** $\lambda(t), t \geq 0$, if $X_0 = 0$ and the increments of X on disjoint intervals are independent Poisson random variables:

$$\mathbf{P}\{X(t) - X(s) = k\} = \frac{\left(\int_s^t \lambda(t) dt\right)^k}{k!} \exp\left\{-\int_s^t \lambda(t) dt\right\}.$$

Observations: n independent realizations $X^{(n)} = (X_1^{(n)}, \dots, X_n^{(n)})$ of a Poisson process with intensity function $\lambda_{\vartheta}^{(n)}(t), 0 \leq t \leq \tau$.

Unknown parameter: $\vartheta \in \Theta = (\alpha, \beta), 0 \leq \alpha < \beta \leq \tau$.

Asymptotics: $n \rightarrow \infty$.



Model of observations

Recall: $X = (X(t), t \geq 0)$ is an (inhomogeneous) **Poisson process** with **intensity function** $\lambda(t), t \geq 0$, if $X_0 = 0$ and the increments of X on disjoint intervals are independent Poisson random variables:

$$\mathbf{P}\{X(t) - X(s) = k\} = \frac{\left(\int_s^t \lambda(t) dt\right)^k}{k!} \exp\left\{-\int_s^t \lambda(t) dt\right\}.$$

Observations: n independent realizations $X^{(n)} = (X_1^{(n)}, \dots, X_n^{(n)})$ of a Poisson process with intensity function $\lambda_{\vartheta}^{(n)}(t), 0 \leq t \leq \tau$.

Unknown parameter: $\vartheta \in \Theta = (\alpha, \beta), 0 \leq \alpha < \beta \leq \tau$.

Asymptotics: $n \rightarrow \infty$.

Remark: equivalent to periodic observation on $[0, n\tau]$ and to one observation on $[0, \tau]$ with intensity function $n\lambda_{\vartheta}^{(n)}(t), 0 \leq t \leq \tau$.



Conditions



Conditions

(C1) The intensity function $\lambda_{\vartheta}^{(n)}(t)$ can be written as $\lambda_{\vartheta}^{(n)}(t) = \psi_n(t) + r_n \mathbb{1}_{\{t > \vartheta\}}$, where ψ_n is continuous on $[0, \tau]$.



Conditions

- (C1)** The intensity function $\lambda_{\vartheta}^{(n)}(t)$ can be written as $\lambda_{\vartheta}^{(n)}(t) = \psi_n(t) + r_n \mathbb{1}_{\{t > \vartheta\}}$, where ψ_n is continuous on $[0, \tau]$.
- (C2)** For all $t \in [0, \tau]$, there exist $\lim_{n \rightarrow +\infty} \psi_n(t) = \psi(t) > 0$ uniformly with respect to t .



Conditions

- (C1) The intensity function $\lambda_{\vartheta}^{(n)}(t)$ can be written as $\lambda_{\vartheta}^{(n)}(t) = \psi_n(t) + r_n \mathbb{1}_{\{t > \vartheta\}}$, where ψ_n is continuous on $[0, \tau]$.
- (C2) For all $t \in [0, \tau]$, there exist $\lim_{n \rightarrow +\infty} \psi_n(t) = \psi(t) > 0$ uniformly with respect to t .
- (C3) We have $r_n \rightarrow r \in \mathbb{R}$. If $r = 0$, we also suppose $n r_n^2 \rightarrow +\infty$.



Conditions

- (C1) The intensity function $\lambda_{\vartheta}^{(n)}(t)$ can be written as $\lambda_{\vartheta}^{(n)}(t) = \psi_n(t) + r_n \mathbb{1}_{\{t > \vartheta\}}$, where ψ_n is continuous on $[0, \tau]$.
- (C2) For all $t \in [0, \tau]$, there exist $\lim_{n \rightarrow +\infty} \psi_n(t) = \psi(t) > 0$ uniformly with respect to t .
- (C3) We have $r_n \rightarrow r \in \mathbb{R}$. If $r = 0$, we also suppose $n r_n^2 \rightarrow +\infty$.
- (C4) There exist some constants $\ell, L > 0$ such that $\ell \leq \lambda_{\vartheta}^{(n)}(t) \leq L$ for all $n \in \mathbb{N}$, $\vartheta \in (\alpha, \beta)$ and $t \in [0, \tau]$.



Conditions

- (C1) The intensity function $\lambda_{\vartheta}^{(n)}(t)$ can be written as $\lambda_{\vartheta}^{(n)}(t) = \psi_n(t) + r_n \mathbb{1}_{\{t > \vartheta\}}$, where ψ_n is continuous on $[0, \tau]$.
- (C2) For all $t \in [0, \tau]$, there exist $\lim_{n \rightarrow +\infty} \psi_n(t) = \psi(t) > 0$ uniformly with respect to t .
- (C3) We have $r_n \rightarrow r \in \mathbb{R}$. If $r = 0$, we also suppose $n r_n^2 \rightarrow +\infty$.
- (C4) There exist some constants $\ell, L > 0$ such that $\ell \leq \lambda_{\vartheta}^{(n)}(t) \leq L$ for all $n \in \mathbb{N}$, $\vartheta \in (\alpha, \beta)$ and $t \in [0, \tau]$.

Remark:

(C1) – (C3) and $r > - \min_{t \in [0, \tau]} \psi(t) \implies$ (C4) for n sufficiently large.



Conditions

- (C1) The intensity function $\lambda_{\vartheta}^{(n)}(t)$ can be written as $\lambda_{\vartheta}^{(n)}(t) = \psi_n(t) + r_n \mathbb{1}_{\{t > \vartheta\}}$, where ψ_n is continuous on $[0, \tau]$.
- (C2) For all $t \in [0, \tau]$, there exist $\lim_{n \rightarrow +\infty} \psi_n(t) = \psi(t) > 0$ uniformly with respect to t .
- (C3) We have $r_n \rightarrow r \in \mathbb{R}$. If $r = 0$, we also suppose $n r_n^2 \rightarrow +\infty$.
- (C4') We have $r > -\min_{t \in [0, \tau]} \psi(t)$.



Conditions

- (C1) The intensity function $\lambda_{\vartheta}^{(n)}(t)$ can be written as $\lambda_{\vartheta}^{(n)}(t) = \psi_n(t) + r_n \mathbb{1}_{\{t > \vartheta\}}$, where ψ_n is continuous on $[0, \tau]$.
- (C2) For all $t \in [0, \tau]$, there exist $\lim_{n \rightarrow +\infty} \psi_n(t) = \psi(t) > 0$ uniformly with respect to t .
- (C3) We have $r_n \rightarrow r \in \mathbb{R}$. If $r = 0$, we also suppose $n r_n^2 \rightarrow +\infty$.
- (C4') We have $r > -\min_{t \in [0, \tau]} \psi(t)$.

Remark:

- (C1) – (C2) \iff (C0) The intensity function $\lambda_{\vartheta}^{(n)}(t)$ can be written as $\lambda_{\vartheta}^{(n)}(t) = \psi(t) + r_n \mathbb{1}_{\{t > \vartheta\}}$, where $\psi > 0$ is continuous on $[0, \tau]$.



Likelihood ratio



Likelihood ratio

The measures corresponding to different values of ϑ are equivalent (and also equivalent to the case $\lambda \equiv 1$).



Likelihood ratio

The measures corresponding to different values of ϑ are equivalent (and also equivalent to the case $\lambda \equiv 1$).

The *likelihood* is a (random) càdlàg function of $\vartheta \in (\alpha, \beta)$:

$$L_n(\vartheta, X^{(n)}) = \exp \left\{ \sum_{j=1}^n \int_{[0, \tau]} \ln \lambda_{\vartheta}^{(n)}(t) \, dX_j^{(n)}(t) - n \int_0^{\tau} [\lambda_{\vartheta}^{(n)}(t) - 1] \, dt \right\}.$$

Likelihood ratio

The measures corresponding to different values of ϑ are equivalent (and also equivalent to the case $\lambda \equiv 1$).

The *likelihood* is a (random) càdlàg function of $\vartheta \in (\alpha, \beta)$:

$$L_n(\vartheta, X^{(n)}) = \exp \left\{ \sum_{j=1}^n \int_{[0, \tau]} \ln \lambda_{\vartheta}^{(n)}(t) \, dX_j^{(n)}(t) - n \int_0^{\tau} [\lambda_{\vartheta}^{(n)}(t) - 1] \, dt \right\}.$$

The *normalized likelihood ratio* is a process with trajectories in $\mathcal{D}_0(\mathbb{R})$:

$$Z_{n, \vartheta}(v) = \frac{L_n(\vartheta_v, X^{(n)})}{L_n(\vartheta, X^{(n)})}, \quad v \in \Theta_n = (\varphi_n^{-1}(\alpha - \vartheta), \varphi_n^{-1}(\beta - \vartheta)),$$

where $\vartheta_v \triangleq \vartheta + v\varphi_n$, and *normalization rate* $\varphi_n = \varphi_n(\vartheta) \searrow 0$ suitably.

Likelihood ratio

The measures corresponding to different values of ϑ are equivalent (and also equivalent to the case $\lambda \equiv 1$).

The *likelihood* is a (random) càdlàg function of $\vartheta \in (\alpha, \beta)$:

$$L_n(\vartheta, X^{(n)}) = \exp \left\{ \sum_{j=1}^n \int_{[0, \tau]} \ln \lambda_{\vartheta}^{(n)}(t) \, dX_j^{(n)}(t) - n \int_0^{\tau} [\lambda_{\vartheta}^{(n)}(t) - 1] \, dt \right\}.$$

The *normalized likelihood ratio* is a process with trajectories in $\mathcal{D}_0(\mathbb{R})$:

$$Z_{n, \vartheta}(v) = \frac{L_n(\vartheta_v, X^{(n)})}{L_n(\vartheta, X^{(n)})}, \quad v \in \Theta_n = (\varphi_n^{-1}(\alpha - \vartheta), \varphi_n^{-1}(\beta - \vartheta)),$$

where $\vartheta_v \triangleq \vartheta + v\varphi_n$, and *normalization rate* $\varphi_n = \begin{cases} \frac{1}{|r|n}, & \text{if } r \neq 0, \\ \frac{\psi(\vartheta)}{n r_n^2}, & \text{if } r = 0. \end{cases}$



The processes Z_ρ and Z_0



The processes Z_ρ and Z_0

The process Z_ρ is a process with trajectories in $\mathcal{D}_0(\mathbb{R})$ defined by

$$Z_\rho(v) = \begin{cases} \exp\{\rho \Pi_+(v) - v\}, & \text{if } v \geq 0, \\ \exp\{-\rho \Pi_-((-v)-) - v\}, & \text{if } v \leq 0, \end{cases}$$

where $\rho > 0$, and Π_+ and Π_- are two independent Poisson processes on \mathbb{R}_+ of intensities $1/(e^\rho - 1)$ and $1/(1 - e^{-\rho})$.



The processes Z_ρ and Z_0

The process Z_ρ is a process with trajectories in $\mathcal{D}_0(\mathbb{R})$ defined by

$$Z_\rho(v) = \begin{cases} \exp\{\rho \Pi_+(v) - v\}, & \text{if } v \geq 0, \\ \exp\{-\rho \Pi_-((-v)-) - v\}, & \text{if } v \leq 0, \end{cases}$$

where $\rho > 0$, and Π_+ and Π_- are two independent Poisson processes on \mathbb{R}_+ of intensities $1/(e^\rho - 1)$ and $1/(1 - e^{-\rho})$.

The process Z_0 is a process with trajectories in $\mathcal{C}_0(\mathbb{R}) \subset \mathcal{D}_0(\mathbb{R})$ defined by

$$Z_0(v) = \exp\left\{W(v) - \frac{|v|}{2}\right\},$$

where W is a standard two-sided Brownian motion (Wiener process).



Limit of the normalized likelihood ratio



Limit of the normalized likelihood ratio

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled. Then, uniformly in ϑ on any compact set $\mathbb{K} \subset \Theta$, the process $Z_{n,\vartheta}$ converges weakly in the space $\mathcal{D}_0(\mathbb{R})$ to



Limit of the normalized likelihood ratio

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled. Then, uniformly in ϑ on any compact set $\mathbb{K} \subset \Theta$, the process $Z_{n,\vartheta}$ converges weakly in the space $\mathcal{D}_0(\mathbb{R})$ to

- ◆ the process Z_ρ with $\rho = \ln \frac{\psi(\vartheta)}{\psi(\vartheta)+r}$, in the case $r < 0$,



Limit of the normalized likelihood ratio

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled. Then, uniformly in ϑ on any compact set $\mathbb{K} \subset \Theta$, the process $Z_{n,\vartheta}$ converges weakly in the space $\mathcal{D}_0(\mathbb{R})$ to

- ◆ the process Z_ρ with $\rho = \ln \frac{\psi(\vartheta)}{\psi(\vartheta)+r}$, in the case $r < 0$,
- ◆ the process Z'_ρ with $\rho = \ln \frac{\psi(\vartheta)+r}{\psi(\vartheta)}$ defined by $Z'_\rho(v) = Z_\rho((-v)-)$, in the case $r > 0$,

Limit of the normalized likelihood ratio

Theorem. — Let the conditions (C1) – (C4) be fulfilled. Then, uniformly in ϑ on any compact set $\mathbb{K} \subset \Theta$, the process $Z_{n,\vartheta}$ converges weakly in the space $\mathcal{D}_0(\mathbb{R})$ to

- ◆ the process Z_ρ with $\rho = \ln \frac{\psi(\vartheta)}{\psi(\vartheta)+r}$, in the case $r < 0$,
- ◆ the process Z'_ρ with $\rho = \ln \frac{\psi(\vartheta)+r}{\psi(\vartheta)}$ defined by $Z'_\rho(v) = Z_\rho((-v)-)$, in the case $r > 0$,
- ◆ the process Z_0 , in the case $r = 0$.



Limit of the normalized likelihood ratio

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled. Then, uniformly in ϑ on any compact set $\mathbb{K} \subset \Theta$, the process $Z_{n,\vartheta}$ converges weakly in the space $\mathcal{D}_0(\mathbb{R})$ to

- ◆ the process Z_ρ with $\rho = \ln \frac{\psi(\vartheta)}{\psi(\vartheta)+r}$, in the case $r < 0$,
- ◆ the process Z'_ρ with $\rho = \ln \frac{\psi(\vartheta)+r}{\psi(\vartheta)}$ defined by
$$Z'_\rho(v) = Z_\rho((-v)-),$$
 in the case $r > 0$,
- ◆ the process Z_0 , in the case $r = 0$.

Remark: in the case $r \neq 0$, the limit is the same as in the case of a fixed jump size ($r_n \equiv r$) studied by Kutoyants (1984, 1998) (see also Dachian, Kutoyants, Yang (2015) for hypotheses testing).



Parameter estimation



Parameter estimation

The *maximum likelihood estimator (MLE)* $\hat{\vartheta}_n$ is defined by

$$\max \left\{ L_n \left(\hat{\vartheta}_n +, X^{(n)} \right), L_n \left(\hat{\vartheta}_n -, X^{(n)} \right) \right\} = \sup_{\vartheta \in (\alpha, \beta)} L_n \left(\vartheta, X^{(n)} \right).$$



Parameter estimation

The *maximum likelihood estimator (MLE)* $\hat{\vartheta}_n$ is defined by

$$\max \left\{ L_n \left(\hat{\vartheta}_n +, X^{(n)} \right), L_n \left(\hat{\vartheta}_n -, X^{(n)} \right) \right\} = \sup_{\vartheta \in (\alpha, \beta)} L_n(\vartheta, X^{(n)}).$$

The *Bayes estimator (BE)* $\tilde{\vartheta}_n$ (for square loss and) for a given *prior density* p is defined by

$$\tilde{\vartheta}_n = \frac{\int_{\alpha}^{\beta} \vartheta p(\vartheta) L_n(\vartheta, X^{(n)}) d\vartheta}{\int_{\alpha}^{\beta} p(\vartheta) L_n(\vartheta, X^{(n)}) d\vartheta}.$$



Asymptotic efficiency



Asymptotic efficiency

We introduce the random variables ξ and ζ by the equations

$$Z_0(\xi) = \sup_{v \in \mathbb{R}} Z_0(v) \quad \text{and} \quad \zeta = \frac{\int_{-\infty}^{+\infty} v Z_0(v) dv}{\int_{-\infty}^{+\infty} Z_0(v) dv}.$$



Asymptotic efficiency

We introduce the random variables ξ and ζ by the equations

$$Z_0(\xi) = \sup_{v \in \mathbb{R}} Z_0(v) \quad \text{and} \quad \zeta = \frac{\int_{-\infty}^{+\infty} v Z_0(v) dv}{\int_{-\infty}^{+\infty} Z_0(v) dv}.$$

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$.

Asymptotic efficiency

We introduce the random variables ξ and ζ by the equations

$$Z_0(\xi) = \sup_{v \in \mathbb{R}} Z_0(v) \quad \text{and} \quad \zeta = \frac{\int_{-\infty}^{+\infty} v Z_0(v) dv}{\int_{-\infty}^{+\infty} Z_0(v) dv}.$$

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$.

Then, for any $\vartheta_0 \in \Theta$, we have

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow +\infty} \inf_{\bar{\vartheta}_n} \sup_{|\vartheta - \vartheta_0| < \delta} \frac{n^2 r_n^4}{\psi^2(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} (\bar{\vartheta}_n - \vartheta)^2 \geq \mathbf{E} \zeta^2,$$

where the inf is taken over all possible estimators $\bar{\vartheta}_n$.

Asymptotic efficiency

We introduce the random variables ξ and ζ by the equations

$$Z_0(\xi) = \sup_{v \in \mathbb{R}} Z_0(v) \quad \text{and} \quad \zeta = \frac{\int_{-\infty}^{+\infty} v Z_0(v) dv}{\int_{-\infty}^{+\infty} Z_0(v) dv}.$$

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$.

Then, for any $\vartheta_0 \in \Theta$, we have

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow +\infty} \inf_{\bar{\vartheta}_n} \sup_{|\vartheta - \vartheta_0| < \delta} \frac{n^2 r_n^4}{\psi^2(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} (\bar{\vartheta}_n - \vartheta)^2 \geq \mathbf{E} \zeta^2 = 16 \zeta(3) \approx 19.23,$$

where the inf is taken over all possible estimators $\bar{\vartheta}_n$.

Asymptotic efficiency

We introduce the random variables ξ and ζ by the equations

$$Z_0(\xi) = \sup_{v \in \mathbb{R}} Z_0(v) \quad \text{and} \quad \zeta = \frac{\int_{-\infty}^{+\infty} v Z_0(v) dv}{\int_{-\infty}^{+\infty} Z_0(v) dv}.$$

Theorem. — Let the conditions (C1) – (C4) be fulfilled with $r = 0$.

Then, for any $\vartheta_0 \in \Theta$, we have

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow +\infty} \inf_{\bar{\vartheta}_n} \sup_{|\vartheta - \vartheta_0| < \delta} \frac{n^2 r_n^4}{\psi^2(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} (\bar{\vartheta}_n - \vartheta)^2 \geq \mathbf{E} \zeta^2 = 16 \zeta(3) \approx 19.23,$$

where the inf is taken over all possible estimators $\bar{\vartheta}_n$.

Definition. — We say that an estimator ϑ_n^* is asymptotically efficient if

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow +\infty} \sup_{|\vartheta - \vartheta_0| < \delta} \frac{n^2 r_n^4}{\psi^2(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} (\vartheta_n^* - \vartheta)^2 = \mathbf{E} \zeta^2 = 16 \zeta(3),$$

for all $\vartheta_0 \in \Theta$.



Properties of the MLE



Properties of the MLE

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$.



Properties of the MLE

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the maximum likelihood estimator $\hat{\vartheta}_n$ satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations



Properties of the MLE

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the maximum likelihood estimator $\hat{\vartheta}_n$ satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

$$\blacklozenge \hat{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta \quad (\text{convergence in probability}),$$



Properties of the MLE

Theorem. — Let the conditions (C1) – (C4) be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the maximum likelihood estimator $\hat{\vartheta}_n$ satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

- ◆ $\hat{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta$ (convergence in probability),
- ◆ $\frac{n r_n^2}{\psi(\vartheta)} (\hat{\vartheta}_n - \vartheta) \Longrightarrow \xi$ (convergence in law),

Properties of the MLE

Theorem. — Let the conditions (C1) – (C4) be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the maximum likelihood estimator $\hat{\vartheta}_n$ satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

- ◆ $\hat{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta$ (convergence in probability),
- ◆ $\frac{n r_n^2}{\psi(\vartheta)} (\hat{\vartheta}_n - \vartheta) \Longrightarrow \xi$ (convergence in law),
- ◆ $\frac{n^p r_n^{2p}}{\psi^p(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} |\hat{\vartheta}_n - \vartheta|^p \longrightarrow \mathbf{E} |\xi|^p$ for any $p > 0$.

Properties of the MLE

Theorem. — Let the conditions (C1) – (C4) be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the maximum likelihood estimator $\hat{\vartheta}_n$ satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

- ◆ $\hat{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta$ (convergence in probability),
- ◆ $\frac{n r_n^2}{\psi(\vartheta)} (\hat{\vartheta}_n - \vartheta) \Longrightarrow \xi$ (convergence in law),
- ◆ $\frac{n^p r_n^{2p}}{\psi^p(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} |\hat{\vartheta}_n - \vartheta|^p \longrightarrow \mathbf{E} |\xi|^p$ for any $p > 0$.

In particular, the relative asymptotic efficiency of $\hat{\vartheta}_n$ is

$$\frac{\mathbf{E} \zeta^2}{\mathbf{E} \xi^2}.$$

Properties of the MLE

Theorem. — Let the conditions (C1) – (C4) be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the maximum likelihood estimator $\hat{\vartheta}_n$ satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

- ◆ $\hat{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta$ (convergence in probability),
- ◆ $\frac{n r_n^2}{\psi(\vartheta)} (\hat{\vartheta}_n - \vartheta) \Longrightarrow \xi$ (convergence in law),
- ◆ $\frac{n^p r_n^{2p}}{\psi^p(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} |\hat{\vartheta}_n - \vartheta|^p \longrightarrow \mathbf{E} |\xi|^p$ for any $p > 0$.

In particular, the relative asymptotic efficiency of $\hat{\vartheta}_n$ is

$$\frac{\mathbf{E}\zeta^2}{\mathbf{E}\xi^2} = \frac{16 \zeta(3)}{26} \approx 0.74.$$



Properties of the BEs



Properties of the BEs

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$.



Properties of the BEs

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the Bayes estimator $\tilde{\vartheta}_n$ for any continuous strictly positive prior density satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations



Properties of the BEs

Theorem. — Let the conditions **(C1)** – **(C4)** be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the Bayes estimator $\tilde{\vartheta}_n$ for any continuous strictly positive prior density satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

$$\blacklozenge \tilde{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta \quad (\text{convergence in probability}),$$



Properties of the BEs

Theorem. — Let the conditions (C1) – (C4) be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the Bayes estimator $\tilde{\vartheta}_n$ for any continuous strictly positive prior density satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

- ◆ $\tilde{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta$ (convergence in probability),
- ◆ $\frac{n r_n^2}{\psi(\vartheta)} (\tilde{\vartheta}_n - \vartheta) \Longrightarrow \zeta$ (convergence in law),



Properties of the BEs

Theorem. — Let the conditions (C1) – (C4) be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the Bayes estimator $\tilde{\vartheta}_n$ for any continuous strictly positive prior density satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

- ◆ $\tilde{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta$ (convergence in probability),
- ◆ $\frac{n r_n^2}{\psi(\vartheta)} (\tilde{\vartheta}_n - \vartheta) \Longrightarrow \zeta$ (convergence in law),
- ◆ $\frac{n^p r_n^{2p}}{\psi^p(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} |\tilde{\vartheta}_n - \vartheta|^p \longrightarrow \mathbf{E} |\zeta|^p$ for any $p > 0$.

Properties of the BEs

Theorem. — Let the conditions (C1) – (C4) be fulfilled with $r = 0$. Then, as $n \rightarrow +\infty$, the Bayes estimator $\tilde{\vartheta}_n$ for any continuous strictly positive prior density satisfies, uniformly on ϑ in any compact set $\mathbb{K} \subset \Theta$, the relations

- ◆ $\tilde{\vartheta}_n \xrightarrow{\mathbf{P}} \vartheta$ (convergence in probability),
- ◆ $\frac{n r_n^2}{\psi(\vartheta)} (\tilde{\vartheta}_n - \vartheta) \Longrightarrow \zeta$ (convergence in law),
- ◆ $\frac{n^p r_n^{2p}}{\psi^p(\vartheta)} \mathbf{E}_{\vartheta}^{(n)} |\tilde{\vartheta}_n - \vartheta|^p \longrightarrow \mathbf{E} |\zeta|^p$ for any $p > 0$.

In particular, $\tilde{\vartheta}_n$ is asymptotically efficient.



Hypothesis testing



Hypothesis testing

Problem: now $\Theta = [\vartheta_0, b)$ and we have to choose between

$$\mathcal{H}_0 \quad : \quad \vartheta = \vartheta_0,$$

$$\mathcal{H}_1 \quad : \quad \vartheta > \vartheta_0.$$



Hypothesis testing

Problem: now $\Theta = [\vartheta_0, b)$ and we have to choose between

$$\mathcal{H}_0 \quad : \quad \vartheta = \vartheta_0,$$

$$\mathcal{H}_1 \quad : \quad \vartheta > \vartheta_0.$$

Tests: a (randomized) **test** $\bar{\psi}_n = \bar{\psi}_n(X^{(n)})$ is defined as the probability to reject the hypothesis \mathcal{H}_0 .

Hypothesis testing

Problem: now $\Theta = [\vartheta_0, b)$ and we have to choose between

$$\mathcal{H}_0 \quad : \quad \vartheta = \vartheta_0,$$

$$\mathcal{H}_1 \quad : \quad \vartheta > \vartheta_0.$$

Tests: a (randomized) **test** $\bar{\psi}_n = \bar{\psi}_n(X^{(n)})$ is defined as the probability to reject the hypothesis \mathcal{H}_0 .

Size: we denote \mathcal{K}_ε the class of tests $\bar{\psi}_n$ of asymptotic **size** $\varepsilon \in [0, 1]$:

$$\mathcal{K}_\varepsilon = \left\{ \bar{\psi}_n \quad : \quad \lim_{n \rightarrow \infty} \mathbf{E}_{\vartheta_0}^{(n)} \bar{\psi}_n(X^{(n)}) = \varepsilon \right\}.$$

Hypothesis testing

Problem: now $\Theta = [\vartheta_0, b)$ and we have to choose between

$$\mathcal{H}_0 \quad : \quad \vartheta = \vartheta_0,$$

$$\mathcal{H}_1 \quad : \quad \vartheta > \vartheta_0.$$

Tests: a (randomized) **test** $\bar{\psi}_n = \bar{\psi}_n(X^{(n)})$ is defined as the probability to reject the hypothesis \mathcal{H}_0 .

Size: we denote \mathcal{K}_ε the class of tests $\bar{\psi}_n$ of asymptotic **size** $\varepsilon \in [0, 1]$:

$$\mathcal{K}_\varepsilon = \left\{ \bar{\psi}_n \quad : \quad \lim_{n \rightarrow \infty} \mathbf{E}_{\vartheta_0}^{(n)} \bar{\psi}_n(X^{(n)}) = \varepsilon \right\}.$$

Power: **power function** of $\bar{\psi}_n$ is $\beta(\bar{\psi}_n, \vartheta) = \mathbf{E}_{\vartheta}^{(n)} \bar{\psi}_n(X^{(n)})$, $\vartheta > \vartheta_0$.



Pitman's approach



Pitman's approach

Close or contiguous alternatives: $\vartheta = \vartheta_u \triangleq \vartheta_0 + u\varphi_n$, where $\varphi_n = \varphi_n(\vartheta_0) \searrow 0$ and $u \in \Theta_n^+ = [0, \varphi_n^{-1}(b - \vartheta_0))$.



Pitman's approach

Close or contiguous alternatives: $\vartheta = \vartheta_u \triangleq \vartheta_0 + u\varphi_n$, where $\varphi_n = \varphi_n(\vartheta_0) \searrow 0$ and $u \in \Theta_n^+ = [0, \varphi_n^{-1}(b - \vartheta_0))$.

Rate: φ_n such that the normalized likelihood ratio

$$Z_{n,\vartheta_0}(v) = \frac{L_n(\vartheta_0 + v\varphi_n, X^{(n)})}{L_n(\vartheta_0, X^{(n)})}, \quad v \in \Theta_n^+,$$

has a non degenerate limit in $\mathcal{D}_0(\mathbb{R}_+)$



Pitman's approach

Close or contiguous alternatives: $\vartheta = \vartheta_u \triangleq \vartheta_0 + u\varphi_n$, where $\varphi_n = \varphi_n(\vartheta_0) \searrow 0$ and $u \in \Theta_n^+ = [0, \varphi_n^{-1}(b - \vartheta_0))$.

Rate: φ_n such that the normalized likelihood ratio

$$Z_{n,\vartheta_0}(v) = \frac{L_n(\vartheta_0 + v\varphi_n, X^{(n)})}{L_n(\vartheta_0, X^{(n)})}, \quad v \in \Theta_n^+,$$

has a non degenerate limit in $\mathcal{D}_0(\mathbb{R}_+)$: $\varphi_n = \frac{\psi(\vartheta_0)}{n r_n^2}$ in our case.

Pitman's approach

Close or contiguous alternatives: $\vartheta = \vartheta_u \triangleq \vartheta_0 + u\varphi_n$, where $\varphi_n = \varphi_n(\vartheta_0) \searrow 0$ and $u \in \Theta_n^+ = [0, \varphi_n^{-1}(b - \vartheta_0))$.

Rate: φ_n such that the normalized likelihood ratio

$$Z_{n,\vartheta_0}(v) = \frac{L_n(\vartheta_0 + v\varphi_n, X^{(n)})}{L_n(\vartheta_0, X^{(n)})}, \quad v \in \Theta_n^+,$$

has a non degenerate limit in $\mathcal{D}_0(\mathbb{R}_+)$: $\varphi_n = \frac{\psi(\vartheta_0)}{n r_n^2}$ in our case.

Problem: we have to choose between

$$\mathcal{H}_0 \quad : \quad u = 0,$$

$$\mathcal{H}_1 \quad : \quad u > 0.$$



Pitman's approach

Close or contiguous alternatives: $\vartheta = \vartheta_u \triangleq \vartheta_0 + u\varphi_n$, where $\varphi_n = \varphi_n(\vartheta_0) \searrow 0$ and $u \in \Theta_n^+ = [0, \varphi_n^{-1}(b - \vartheta_0))$.

Rate: φ_n such that the normalized likelihood ratio

$$Z_{n,\vartheta_0}(v) = \frac{L_n(\vartheta_0 + v\varphi_n, X^{(n)})}{L_n(\vartheta_0, X^{(n)})}, \quad v \in \Theta_n^+,$$

has a non degenerate limit in $\mathcal{D}_0(\mathbb{R}_+)$: $\varphi_n = \frac{\psi(\vartheta_0)}{n r_n^2}$ in our case.

Problem: we have to choose between

$$\mathcal{H}_0 \quad : \quad u = 0,$$

$$\mathcal{H}_1 \quad : \quad u > 0.$$

Power: power function of $\bar{\psi}_n$ is $\beta(\bar{\psi}_n, u) = \mathbf{E}_{\vartheta_u}^{(n)} \bar{\psi}_n(X^{(n)}), u > 0$.



Limits of the likelihood ratio



Limits of the likelihood ratio

Limit under hypothesis: under $\vartheta = \vartheta_0$, the limit of the normalized likelihood ratio Z_{n,ϑ_0} is

$$Z_0(v) = \exp\left\{W(v) - \frac{v}{2}\right\}, \quad v \geq 0,$$

where W is a standard Brownian motion (Wiener process).



Limits of the likelihood ratio

Limit under hypothesis: under $\vartheta = \vartheta_0$, the limit of the normalized likelihood ratio Z_{n,ϑ_0} is

$$Z_0(v) = \exp\left\{W(v) - \frac{v}{2}\right\}, \quad v \geq 0,$$

where W is a standard Brownian motion (Wiener process).

Limit under alternative: under $\vartheta = \vartheta_u$ (with any fixed $u > 0$), the limit of the normalized likelihood ratio Z_{n,ϑ_0} is

$$Z_0^{(u)}(v) = \exp\left\{W(v) - \frac{|v - u|}{2} + \frac{u}{2}\right\}, \quad v \geq 0,$$

where W is a standard Brownian motion (Wiener process).



Wald's test



Wald's test

The **maximum likelihood estimator (MLE)** $\hat{\vartheta}_n$ is now given by

$$\max \left\{ L_n(\hat{\vartheta}_n +, X^{(n)}), L_n(\hat{\vartheta}_n -, X^{(n)}) \right\} = \sup_{\vartheta \in [\vartheta_0, b)} L_n(\vartheta, X^{(n)}).$$



Wald's test

The **maximum likelihood estimator (MLE)** $\hat{\vartheta}_n$ is now given by

$$\max \left\{ L_n(\hat{\vartheta}_n +, X^{(n)}), L_n(\hat{\vartheta}_n -, X^{(n)}) \right\} = \sup_{\vartheta \in [\vartheta_0, b)} L_n(\vartheta, X^{(n)}).$$

The **Wald's test (WT)** is based on the MLE $\hat{\vartheta}_n$ and is defined by

$$\phi_n^\circ(X^{(n)}) = \mathbb{1}_{\left\{ \frac{n r_n^2}{\psi(\vartheta_0)} (\hat{\vartheta}_n - \vartheta_0) > m_\varepsilon \right\}}$$

with m_ε solution of

$$\int_{m_\varepsilon}^{+\infty} \left(\frac{1}{\sqrt{2\pi t}} \exp\{-t/8\} - \frac{1}{2} \Phi(-\sqrt{t}/2) \right) dt = \varepsilon,$$

where Φ is the distribution function of $\mathcal{N}(0, 1)$.



Properties of the WT



Properties of the WT

- ◆ The test ϕ_n° belongs to \mathcal{K}_ε .



Properties of the WT

- ◆ The test ϕ_n° belongs to \mathcal{K}_ε .
- ◆ The power function of ϕ_n° converges to $\beta^\circ(u)$:

$$\beta(\phi_n^\circ, u) \longrightarrow \beta^\circ(u),$$

where

$$\beta^\circ(u) = \mathbf{P} \{ \xi_u > m_\varepsilon - u \}, \quad Z_0(\xi_u) = \sup_{v \geq -u} Z_0(v)$$

and/or

$$\beta^\circ(u) = \mathbf{P} \{ \xi_+^{(u)} > m_\varepsilon \}, \quad Z_0^{(u)}(\xi_+^{(u)}) = \sup_{v \geq 0} Z_0^{(u)}(v).$$



General likelihood ratio test



General likelihood ratio test

The **general likelihood ratio test (GLRT)** is defined by

$$\hat{\phi}_n(X^{(n)}) = \mathbb{1}_{\{Q(X^{(n)}) > 1/\varepsilon\}}$$

with

$$\begin{aligned} Q(X^{(n)}) &= \sup_{\vartheta \in [\vartheta_0, b)} \frac{L_n(\vartheta, X^{(n)})}{L_n(\vartheta_0, X^{(n)})} \\ &= \frac{\max\left\{L_n(\hat{\vartheta}_n^+, X^{(n)}), L_n(\hat{\vartheta}_n^-, X^{(n)})\right\}}{L_n(\vartheta_0, X^{(n)})}. \end{aligned}$$



Properties of the GLRT



Properties of the GLRT

- ◆ The test $\hat{\phi}_n$ belongs to \mathcal{K}_ε .



Properties of the GLRT

- ◆ The test $\hat{\phi}_n$ belongs to \mathcal{K}_ε .
- ◆ The power function of $\hat{\phi}_n$ converges to $\hat{\beta}(u)$:

$$\beta(\hat{\psi}_n, u) \longrightarrow \hat{\beta}(u),$$

where

$$\hat{\beta}(u) = \mathbf{P}\{\hat{Z}_u > 1/\varepsilon\}, \quad \hat{Z}_u = \left(Z_0(-u)\right)^{-1} \sup_{v \geq -u} Z_0(v)$$

and/or

$$\hat{\beta}(u) = \mathbf{P}\{\hat{Z}_+^{(u)} > 1/\varepsilon\}, \quad \hat{Z}_+^{(u)} = \sup_{v \geq 0} Z_0^{(u)}(v).$$



First Bayes test



First Bayes test

The *Bayes estimator (BE)* $\tilde{\vartheta}_n$ (for square loss and) for a given *prior density* p is now given by

$$\tilde{\vartheta}_n = \frac{\int_{\vartheta_0}^b \vartheta p(\vartheta) L_n(\vartheta, X^{(n)}) d\vartheta}{\int_{\vartheta_0}^b p(\vartheta) L_n(\vartheta, X^{(n)}) d\vartheta}.$$

First Bayes test

The *Bayes estimator (BE)* $\tilde{\vartheta}_n$ (for square loss and) for a given *prior density* p is now given by

$$\tilde{\vartheta}_n = \frac{\int_{\vartheta_0}^b \vartheta p(\vartheta) L_n(\vartheta, X^{(n)}) d\vartheta}{\int_{\vartheta_0}^b p(\vartheta) L_n(\vartheta, X^{(n)}) d\vartheta}.$$

The first **Bayes test (BT1)** is a Wald-type test based on the BE $\tilde{\vartheta}_n$ and is defined by

$$\tilde{\phi}_n(X^{(n)}) = \mathbb{1}_{\left\{ \frac{n r_n^2}{\psi(\vartheta_0)} (\tilde{\vartheta}_n - \vartheta_0) > k_\varepsilon \right\}}$$

with k_ε solution of

$$\mathbf{P} \{ \zeta_+ > k_\varepsilon \} = \varepsilon, \quad \zeta_+ = \frac{\int_0^{+\infty} v Z_0(v) dv}{\int_0^{+\infty} Z_0(v) dv}.$$



Properties of the BT1



Properties of the BT1

- ◆ The test $\tilde{\phi}_n$ belongs to \mathcal{K}_ε .

Properties of the BT1

- ◆ The test $\tilde{\phi}_n$ belongs to \mathcal{K}_ε .
- ◆ The power function of $\tilde{\phi}_n$ converges to $\tilde{\beta}(u)$:

$$\beta(\tilde{\phi}_n, u) \longrightarrow \tilde{\beta}(u),$$

where

$$\tilde{\beta}(u) = \mathbf{P} \{ \zeta_u > k_\varepsilon - u \}, \quad \zeta_u = \frac{\int_{-u}^{+\infty} v Z_0(v) \, dv}{\int_{-u}^{+\infty} Z_0(v) \, dv}$$

and/or

$$\tilde{\beta}(u) = \mathbf{P} \{ \zeta_+^{(u)} > k_\varepsilon \}, \quad \zeta_+^{(u)} = \frac{\int_0^{+\infty} v Z_0^{(u)}(v) \, dv}{\int_0^{+\infty} Z_0^{(u)}(v) \, dv}.$$



Second Bayes test



Second Bayes test

For a test $\bar{\phi}_n$, the **mean (or averaged) power** is

$$\beta(\bar{\phi}_n) = \int_{\vartheta_0}^b \beta(\bar{\phi}_n, \vartheta) p(\vartheta) d\vartheta.$$

Second Bayes test

For a test $\bar{\phi}_n$, the **mean (or averaged) power** is

$$\beta(\bar{\phi}_n) = \int_{\vartheta_0}^b \beta(\bar{\phi}_n, \vartheta) p(\vartheta) d\vartheta.$$

The second **Bayes test (BT2)** is the test which maximizes the mean power and is defined by

$$\tilde{\phi}_n^*(X^{(n)}) = \mathbb{1}_{\{R(X^{(n)}) > g_\varepsilon\}}$$

with

$$R(X^{(n)}) = \frac{n r_n^2 \int_{\vartheta_0}^b L_n(\vartheta, X^{(n)}) p(\vartheta) d\vartheta}{p(\vartheta_0) \psi(\vartheta_0) L_n(\vartheta_0, X^{(n)})}$$

and g_ε solution of

$$\mathbf{P}\{\tilde{Z}_+ > g_\varepsilon\} = \varepsilon, \quad \tilde{Z}_+ = \int_0^{+\infty} Z_0(v) dv.$$



Properties of the BT2



Properties of the BT2

- ◆ The test $\tilde{\phi}_n^*$ belongs to \mathcal{K}_ε .

Properties of the BT2

- ◆ The test $\tilde{\phi}_n^*$ belongs to \mathcal{K}_ε .
- ◆ The power function of $\tilde{\phi}_n^*$ converges to $\tilde{\beta}^*(u)$:

$$\beta(\tilde{\phi}_n^*, u) \longrightarrow \tilde{\beta}^*(u),$$

where

$$\tilde{\beta}^*(u) = \mathbf{P}\{\tilde{Z}_u > g_\varepsilon\}, \quad \tilde{Z}_u = (Z_0(-u))^{-1} \int_{-u}^{+\infty} Z_0(v) \, dv$$

and/or

$$\tilde{\beta}^*(u) = \mathbf{P}\{\tilde{Z}_+^{(u)} > g_\varepsilon\}, \quad \tilde{Z}_+^{(u)} = \int_0^{+\infty} Z_0^{(u)}(v) \, dv.$$



Numerical simulations



Numerical simulations

◆ $\lambda_{\vartheta}^{(n)}(t) = 1.5 + n^{-0.25} \mathbb{1}_{\{t > \vartheta\}}, \quad 0 \leq t \leq 4.$



Numerical simulations

- ◆ $\lambda_{\vartheta}^{(n)}(t) = 1.5 + n^{-0.25} \mathbb{1}_{\{t > \vartheta\}}, \quad 0 \leq t \leq 4.$
- ◆ $\vartheta_0 = 2$ and $\Theta = [2, 4).$



Numerical simulations

- ◆ $\lambda_{\vartheta}^{(n)}(t) = 1.5 + n^{-0.25} \mathbb{1}_{\{t > \vartheta\}}, \quad 0 \leq t \leq 4.$
- ◆ $\vartheta_0 = 2$ and $\Theta = [2, 4).$
- ◆ $\varphi_n = \frac{\psi(\vartheta_0)}{n r_n^2} = \frac{1.5}{\sqrt{n}}.$



Numerical simulations

- ◆ $\lambda_{\vartheta}^{(n)}(t) = 1.5 + n^{-0.25} \mathbb{1}_{\{t > \vartheta\}}, \quad 0 \leq t \leq 4.$
- ◆ $\vartheta_0 = 2$ and $\Theta = [2, 4).$
- ◆ $\varphi_n = \frac{\psi(\vartheta_0)}{n r_n^2} = \frac{1.5}{\sqrt{n}}.$
- ◆ prior distribution for BT1: uniform on $[2, 4).$

Numerical simulations





- ◆ $\lambda_{\vartheta}^{(n)}(t) = 1.5 + n^{-0.25} \mathbb{1}_{\{t > \vartheta\}}, \quad 0 \leq t \leq 4.$
- ◆ $\vartheta_0 = 2$ and $\Theta = [2, 4).$
- ◆ $\varphi_n = \frac{\psi(\vartheta_0)}{n r_n^2} = \frac{1.5}{\sqrt{n}}.$
- ◆ prior distribution for BT1: uniform on $[2, 4).$
- ◆ some thresholds for WT and BT1:

ε	0.001	0.005	0.01	0.05	0.1	0.2
m_{ε}	30.336	20.686	14.886	7.282	4.531	2.236
k_{ε}	24.877	17.588	16.782	8.582	5.573	3.024




Likelihood ratio

Likelihood ratio



Likelihood ratio




Some realization of Z_{n,ϑ_0}




Convergence of power functions

Convergence of power functions



Convergence of power functions




Convergence of power functions of the GLRT, of the WT and of the BT1




Comparison of limit power functions

Comparison of limit power functions



Comparison of limit power functions



Comparison of limit power functions with the Neyman-Pearson envelope



References



References

- ◆ Dachian S. and Yang L., “*On a Poissonian change-point model with variable jump size*”, Statistical Inference for Stochastic Processes **18**, pp. 127–150, 2015.



References

- ◆ Dachian S. and Yang L., “*On a Poissonian change-point model with variable jump size*”, *Statistical Inference for Stochastic Processes* **18**, pp. 127–150, 2015.
- ◆ Kutoyants, Yu. A., “*Parameter Estimation for Stochastic Processes*”, Heldermann-Verlag, Berlin, 1984.
- ◆ Kutoyants, Yu. A., “*Statistical Inference for Spatial Poisson Processes*”, *Lecture Notes in Statistics* **134**, Springer-Verlag, New York, 1998.
- ◆ Dachian S., Kutoyants Yu.A. and Yang L., “*On hypothesis testing for Poisson processes. Singular cases*”, *Communications in Statistics — Theory and Methods* **45**, pp. 6833–6859, 2016.