

Affine Volterra processes and their characteristic functions

Martin Larsson

Department of Mathematics, ETH Zurich

(joint work with Eduardo Abi Jaber and Sergio Pulido)

Asymptotic Statistics of Stochastic Processes and Applications

July 17–21, 2017, Peterhof, Russia

Rough volatility models

- ▶ Empirical studies indicate **volatility is rougher than BM**: Gatheral, Jaisson & Rosenbaum ('14); Bennedsen, Lunde, Pakkanen ('16), ...
- ▶ Subsequent development of **stochastic models with this feature**: Gatheral, Jaisson & Rosenbaum ('14); Guennoun, Jacquier & Roome ('14); Bayer, Friz & Gatheral (15); Bennedsen, Lunde, Pakkanen ('16); **El Euch & Rosenbaum ('16,'17)**, ...
- ▶ These models are able to
 - match roughness of time series data
 - fit implied volatility smiles remarkably well
 - admit in some cases microstructural justification
- ▶ Mathematically, this rests on **fractional Brownian motion** in the tradition of Kolmogorov ('40), Mandelbrot & van Ness ('68), ...

Rough Heston model

- The **Heston model** is the stock price model

$$\frac{dS_t}{S_t} = \sqrt{X_t} d\widetilde{W}_t$$

where the volatility follows a CIR process,

$$X_t = X_0 + \int_0^t \kappa(\theta - X_s) ds + \int_0^t \sigma \sqrt{X_s} dW_s$$

Rough Heston model

- The **Heston model** is the stock price model

$$\frac{dS_t}{S_t} = \sqrt{X_t} d\widetilde{W}_t$$

where the volatility follows a CIR process,

$$X_t = X_0 + \int_0^t \kappa(\theta - X_s) ds + \int_0^t \sigma \sqrt{X_s} dW_s$$

- **El Euch & Rosenbaum ('16)** study the **rough Heston model** obtained by replacing the CIR process by the **rough CIR process**

$$X_t = X_0 + \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} \left(\kappa(\theta - X_s) ds + \sigma \sqrt{X_s} dW_s \right)$$

where $\alpha \in (\frac{1}{2}, 1)$.

Rough Heston model

- ▶ Inspired by the Riemann–Liouville fractional Brownian motion introduced by Lévy ('53)
- ▶ Hölder continuous paths of any order less than $H = \alpha - \frac{1}{2}$
- ▶ Microstructural foundation as scaling limit of Hawkes processes

Rough Heston model

- ▶ Inspired by the Riemann–Liouville fractional Brownian motion introduced by Lévy ('53)
- ▶ Hölder continuous paths of any order less than $H = \alpha - \frac{1}{2}$
- ▶ Microstructural foundation as scaling limit of Hawkes processes

But:

- ▶ Existence and uniqueness is non-trivial: El Euch & Rosenbaum construct the rough CIR using Hawkes processes.
- ▶ Not a semimartingale, not Markovian . . .
- ▶ . . . so how to price options? This is needed for implied volatilities!

Rough Heston model

- ▶ Inspired by the Riemann–Liouville fractional Brownian motion introduced by Lévy ('53)
- ▶ Hölder continuous paths of any order less than $H = \alpha - \frac{1}{2}$
- ▶ Microstructural foundation as scaling limit of Hawkes processes

But:

- ▶ Existence and uniqueness is non-trivial: El Euch & Rosenbaum construct the rough CIR using Hawkes processes.
- ▶ Not a semimartingale, not Markovian . . .
- ▶ . . . so how to price options? This is needed for implied volatilities!

Warmup: The standard Heston model

Characteristic function of the (standard) Heston model

- The Heston model is tractable because $(\log S_t, X_t)$ **is affine**. This gives **explicit characteristic function**:

$$\mathbb{E}[e^{u \log S_T}] = e^{\phi(T) + \psi(T) X_0}$$

for $u \in \mathbb{R}$ and $S_0 = 1$, where (ϕ, ψ) solves the **Riccati equations**

$$\phi' = \kappa\theta\psi \qquad \phi(0) = 0$$

$$\psi' = \frac{1}{2}(u^2 - u) - (u\rho\sigma - \kappa)\psi + \frac{\sigma^2}{2}\psi^2 \qquad \psi(0) = 0$$

Characteristic function of the (standard) Heston model

- The Heston model is tractable because $(\log S_t, X_t)$ **is affine**. This gives **explicit characteristic function**:

$$\mathbb{E}[e^{u \log S_T}] = e^{\phi(T) + \psi(T) X_0}$$

for $u \in \mathbb{R}$ and $S_0 = 1$, where (ϕ, ψ) solves the **Riccati equations**

$$\phi' = \kappa\theta\psi \qquad \phi(0) = 0$$

$$\psi' = \frac{1}{2}(u^2 - u) - (u\rho\sigma - \kappa)\psi + \frac{\sigma^2}{2}\psi^2 \qquad \psi(0) = 0$$

Proof: Define $M_t = e^{\phi(T-t) + \psi(T-t)X_t + u \log S_t}$.

Characteristic function of the (standard) Heston model

- The Heston model is tractable because $(\log S_t, X_t)$ is **affine**. This gives **explicit characteristic function**:

$$\mathbb{E}[e^{u \log S_T}] = e^{\phi(T) + \psi(T) X_0}$$

for $u \in \mathbb{R}$ and $S_0 = 1$, where (ϕ, ψ) solves the **Riccati equations**

$$\phi' = \kappa \theta \psi \qquad \phi(0) = 0$$

$$\psi' = \frac{1}{2}(u^2 - u) - (u\rho\sigma - \kappa)\psi + \frac{\sigma^2}{2}\psi^2 \qquad \psi(0) = 0$$

Proof: Define $M_t = e^{\phi(T-t) + \psi(T-t)X_t + u \log S_t}$. Apply Itô:

$$\frac{dM_t}{M_t} = - \left\{ (\phi' - \kappa \theta \psi) + \left(\psi' - \left[\frac{1}{2}(u^2 - u) - \dots \right] \right) X_t \right\} dt + (dW_t \text{ term})$$

Characteristic function of the (standard) Heston model

- The Heston model is tractable because $(\log S_t, X_t)$ **is affine**. This gives **explicit characteristic function**:

$$\mathbb{E}[e^{u \log S_T}] = e^{\phi(T) + \psi(T) X_0}$$

for $u \in \mathbb{R}$ and $S_0 = 1$, where (ϕ, ψ) solves the **Riccati equations**

$$\phi' = \kappa\theta\psi \qquad \phi(0) = 0$$

$$\psi' = \frac{1}{2}(u^2 - u) - (u\rho\sigma - \kappa)\psi + \frac{\sigma^2}{2}\psi^2 \qquad \psi(0) = 0$$

Proof: Define $M_t = e^{\phi(T-t) + \psi(T-t)X_t + u \log S_t}$. Apply Itô:

$$\frac{dM_t}{M_t} = - \left\{ (\phi' - \kappa\theta\psi) + \left(\psi' - \left[\frac{1}{2}(u^2 - u) - \dots \right] \right) X_t \right\} dt + (dW_t \text{ term})$$

Hence, provided M is a martingale (which follows from $\operatorname{Re} \psi \leq 0$),

$$\mathbb{E}[e^{u \log S_T}] = \mathbb{E}[M_T] = M_0 = e^{\phi(T) + \psi(T) X_0}.$$

What about the rough Heston model?

- ▶ Remarkably, El Euch & Rosenbaum obtain an analogous result for the rough Heston model.
- ▶ **Notation:** $D^\alpha h(t) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t (t-s)^{-\alpha} h(s) ds$

What about the rough Heston model?

- ▶ Remarkably, El Euch & Rosenbaum obtain an analogous result for the rough Heston model.
- ▶ **Notation:** $D^\alpha h(t) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t (t-s)^{-\alpha} h(s) ds$

Theorem. (El Euch & Rosenbaum, '16) Assume we are given a solution ψ of the “**fractional Riccati equation**”

$$D^\alpha \psi = \frac{1}{2}(u^2 - u) + (u\rho\sigma - \kappa)\psi + \frac{\sigma^2}{2}\psi^2$$

and define ϕ and χ by

$$\phi' = \kappa\theta\chi, \quad \phi(0) = 0; \quad \chi' = D^\alpha \psi, \quad \chi(0) = 0.$$

Then

$$\mathbb{E}[e^{u \log S_T}] = e^{\phi(T) + \chi(T)X_0}$$

What about the rough Heston model?

- ▶ Remarkably, El Euch & Rosenbaum obtain an analogous result for the rough Heston model.

- ▶ **Notation:** $D^\alpha h(t) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t (t-s)^{-\alpha} h(s) ds$

Theorem. (El Euch & Rosenbaum, '16) Assume we are given a solution ψ of the “**fractional Riccati equation**”

$$D^\alpha \psi = \frac{1}{2}(u^2 - u) + (u\rho\sigma - \kappa)\psi + \frac{\sigma^2}{2}\psi^2$$

and define ϕ and χ by

$$\phi' = \kappa\theta\chi, \quad \phi(0) = 0; \quad \chi' = D^\alpha \psi, \quad \chi(0) = 0.$$

Then

$$\mathbb{E}[e^{u \log S_T}] = e^{\phi(T) + \chi(T)X_0}$$

Proof: Rather involved. Uses the Hawkes approximation.

Questions

- ▶ Can the proof of this result be simplified?
- ▶ Can the Hawkes approximation be avoided?
- ▶ What about the joint characteristic function of $(\log S_T, X_T)$?
- ▶ What about conditional characteristic function?
- ▶ More general specifications?

Affine Volterra processes

- ▶ State space $E \subseteq \mathbb{R}^d$
- ▶ Affine diffusion and drift coefficients

$$a(x) = A^0 + A^1 x_1 + \cdots + A^d x_d$$

$$b(x) = b^0 + b^1 x_1 + \cdots + b^d x_d$$

with $A^i \in \mathbb{S}^d$, $b^i \in \mathbb{R}^d$, and $a(x) \succeq 0$ on E .

- ▶ $\sigma: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$ continuous with $\sigma(x)\sigma(x)^\top = a(x)$ on E .
- ▶ Matrix-valued kernel $K \in L^2_{\text{loc}}(\mathbb{R}_+, \mathbb{R}^{d \times d})$.

A continuous E -valued solution X of the stochastic Volterra equation

$$X_t = X_0 + \int_0^t K(t-s)b(X_s)ds + \int_0^t K(t-s)\sigma(X_s)dW_s$$

is called an **affine Volterra process** (of convolution type).

Affine Volterra processes

$$X_t = X_0 + \int_0^t K(t-s)b(X_s)ds + \int_0^t K(t-s)\sigma(X_s)dW_s$$

- ▶ **Example:** For usual affine diffusions, take $K(t) \equiv \text{id}$.
- ▶ **Example:** The volatility process in the rough volatility model by Rosenbaum & El Euch is obtained with

$$K(t) = \frac{1}{\Gamma(\alpha)} t^{\alpha-1}$$

- ▶ **Example:** More generally, the full rough volatility model uses $d = 2$ and the kernel

$$K(t) = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{\Gamma(\alpha)} t^{\alpha-1} \end{pmatrix}$$

Characteristic function

Notation: $(F * G)(t) = \int_0^t F(t-s)G(s)ds.$

Characteristic function

Notation: $(F * G)(t) = \int_0^t F(t-s)G(s)ds.$

Theorem. Let $d = 1$. Fix $u \in \mathbb{C}$, assume $\psi \in L^2_{\text{loc}}(\mathbb{R}_+, \mathbb{C})$ solves

$$\psi = uK + \left(\psi b^1 + \frac{A^1}{2} \psi^2 \right) * K, \quad (1)$$

and define ϕ and χ by $\phi(0) = 0$, $\chi(0) = u$, and

$$\phi' = \psi b^0 + \frac{A^0}{2} \psi^2, \quad \chi' = \psi b^1 + \frac{A^1}{2} \psi^2. \quad (2)$$

Then, under a martingale condition,

$$\mathbb{E}[e^{uX_T}] = e^{\phi(T) + \chi(T)X_0}, \quad T \geq 0.$$

Characteristic function

Proof.

- **Ansatz:** Fix T and consider the semimartingale $M_t = e^{Y_t}$, where

$$Y_t = \phi(T - t) + \chi(T)X_0 - \int_0^t \chi'(T - s)X_s ds + \int_0^t \psi(T - s)dZ_s$$

with $Z_t = \int_0^t b(X_s)ds + \int_0^t \sigma(X_s)dW_s$.

Characteristic function

Proof.

- **Ansatz:** Fix T and consider the semimartingale $M_t = e^{Y_t}$, where

$$Y_t = \phi(T-t) + \chi(T)X_0 - \int_0^t \chi'(T-s)X_s ds + \int_0^t \psi(T-s)dZ_s$$

with $Z_t = \int_0^t b(X_s)ds + \int_0^t \sigma(X_s)dW_s$.

- **Itô** yields

$$\begin{aligned} \frac{dM_t}{M_t} &= \left(-\phi' + \psi b^0 + \frac{A^0}{2} \psi^2 \right) dt \\ &\quad + \left(-\chi' + \psi b^1 + \frac{A^1}{2} \psi^2 \right) X_t dt + (dW_t \text{ term}), \end{aligned}$$

a local martingale **by (2)**. By “martingale condition”, a martingale.

Characteristic function

Proof.

- **Ansatz:** Fix T and consider the semimartingale $M_t = e^{Y_t}$, where

$$Y_t = \phi(T-t) + \chi(T)X_0 - \int_0^t \chi'(T-s)X_s ds + \int_0^t \psi(T-s)dZ_s$$

with $Z_t = \int_0^t b(X_s)ds + \int_0^t \sigma(X_s)dW_s$.

- **Itô** yields

$$\begin{aligned} \frac{dM_t}{M_t} = & \left(-\phi' + \psi b^0 + \frac{A^0}{2} \psi^2 \right) dt \\ & + \left(-\chi' + \psi b^1 + \frac{A^1}{2} \psi^2 \right) X_t dt + (dW_t \text{ term}), \end{aligned}$$

a local martingale **by (2)**. By “martingale condition”, a martingale.

- **(1) and stochastic Fubini** yield $Y_T = \phi(0) + \chi(0)X_T$.

Characteristic function

Proof.

- **Ansatz:** Fix T and consider the semimartingale $M_t = e^{Y_t}$, where

$$Y_t = \phi(T-t) + \chi(T)X_0 - \int_0^t \chi'(T-s)X_s ds + \int_0^t \psi(T-s)dZ_s$$

with $Z_t = \int_0^t b(X_s)ds + \int_0^t \sigma(X_s)dW_s$.

- **Itô** yields

$$\begin{aligned} \frac{dM_t}{M_t} &= \left(-\phi' + \psi b^0 + \frac{A^0}{2} \psi^2 \right) dt \\ &\quad + \left(-\chi' + \psi b^1 + \frac{A^1}{2} \psi^2 \right) X_t dt + (dW_t \text{ term}), \end{aligned}$$

a local martingale **by (2)**. By “martingale condition”, a martingale.

- **(1) and stochastic Fubini** yield $Y_T = \phi(0) + \chi(0)X_T$.
- Hence $\mathbb{E}[e^{uX_T}] = \mathbb{E}[M_T] = M_0 = e^{\phi(T) + \chi(T)X_0}$

Characteristic function

- ▶ The above theorem is **probably the quickest way to derive the characteristic function** of $X_t \dots$
- ▶ \dots although the “**martingale condition**” needed to make the derivation rigorous requires additional work.
- ▶ However, the computations do not generalize to **conditional characteristic functions**, which is a nontrivial extension due to lack of Markovian structure.
- ▶ Different, more general, and more informative results are possible.

Conditional expectation

- Set $B = [b^1 \ b^2 \ \dots \ b^d] \in \mathbb{R}^{d \times d}$ and let temporarily $b^0 = 0$:

$$X_t = X_0 + \int_0^t K(t-s) B X_s ds + \int_0^t K(t-s) \sigma(X_s) dW_s$$

Conditional expectation

- Set $B = [b^1 \ b^2 \ \dots \ b^d] \in \mathbb{R}^{d \times d}$ and let temporarily $b^0 = 0$:

$$X_t = X_0 + \int_0^t K(t-s) B X_s ds + \int_0^t K(t-s) \sigma(X_s) dW_s$$

- The kernel $-KB$ admits a **resolvent** $R_B \in L_{\text{loc}}^2(\mathbb{R}_+, \mathbb{R}^{d \times d})$:

$$(KB) * R_B = R_B * (KB) = KB + R_B$$

Conditional expectation

- ▶ Set $B = [b^1 \ b^2 \ \dots \ b^d] \in \mathbb{R}^{d \times d}$ and let temporarily $b^0 = 0$:

$$X_t = X_0 + \int_0^t K(t-s)BX_s ds + \int_0^t K(t-s)\sigma(X_s)dW_s$$

- ▶ The kernel $-KB$ admits a **resolvent** $R_B \in L_{\text{loc}}^2(\mathbb{R}_+, \mathbb{R}^{d \times d})$:

$$(KB) * R_B = R_B * (KB) = KB + R_B$$

- ▶ **Example:** If $K \equiv \text{id}$ then $R_B(t) = -Be^{tB}$.
- ▶ **Example:** If $K(t) = \frac{1}{\Gamma(\alpha)}t^{\alpha-1}$ and $B = -\kappa < 0$, then

$$R_B = f^{\alpha, \kappa}$$

is the so-called **Mittag-Leffler density function**.

Conditional expectation

Theorem. The conditional mean, or **forward process**, is given by

$$\begin{aligned}\mathbb{E}[X_T \mid \mathcal{F}_t] = & \left(\text{id} - \int_0^T R_B(s) ds \right) X_0 + \left(\int_0^T E_B(s) ds \right) b^0 \\ & + \int_0^t E_B(T-s) \sigma(X_s) dW_s,\end{aligned}$$

where R_B is the resolvent of $-KB$ and $E_B = K - R_B * K$.

Conditional characteristic function

► **Recap:**

$$X_t = X_0 + \int_0^t K(t-s)b(X_s)ds + \int_0^t K(t-s)\sigma(X_s)dW_s$$

with

$$a(x) = A^0 + A^1x_1 + \cdots + A^dx_d = \sigma(x)\sigma(x)^\top$$

$$b(x) = b^0 + Bx$$

- **Notation:** For any row vector $u \in (\mathbb{C}^d)^*$ we define the row vector

$$A(u) = (uA^1u^\top, \dots, uA^du^\top).$$

- **Riccati–Volterra equation:** $\psi \in L^2_{\text{loc}}(\mathbb{R}_+, (\mathbb{C}^d)^*)$ such that

$$\psi = uK + \left(\psi B + \frac{1}{2}A(\psi) \right) * K.$$

Conditional characteristic function

Theorem (*). Let $u \in (\mathbb{C}^d)^*$ and assume the Riccati–Volterra equation has solution ψ . Fix $T < \infty$ and define

$$dY_t = \psi(T-t)\sigma(X_t)dW_t - \frac{1}{2}\psi(T-t)a(X_t)\psi(T-t)^\top dt,$$
$$Y_0 = uX_0 + \int_0^T \left(\psi(s)b(X_0) + \frac{1}{2}\psi(s)a(X_0)\psi(s)^\top \right) ds.$$

Then for $t \leq T$,

$$Y_t = \mathbb{E}[uX_T \mid \mathcal{F}_t] + \frac{1}{2} \int_t^T \psi(T-s)a(\mathbb{E}[X_s \mid \mathcal{F}_t])\psi(T-s)^\top ds.$$

Consequently $\{\exp(Y_t), 0 \leq t \leq T\}$ is a local martingale and, if it is a true martingale, one has the affine transform formula

$$\mathbb{E}[e^{uX_T} \mid \mathcal{F}_t] = e^{Y_t}, \quad t \leq T.$$

Conditional characteristic function

Remark. With the same method we actually get a formula for

$$\mathbb{E} \left[\exp \left(u X_T + (f * X)_T \right) \mid \mathcal{F}_t \right]$$

for $u \in (\mathbb{C}^d)^*$ and $f \in L^1_{\text{loc}}(\mathbb{R}_+, (\mathbb{C}^d)^*)$.

Volterra–Ornstein–Uhlenbeck process

- ▶ With $E = \mathbb{R}^d$ and $\sigma(x) \equiv \sigma$ constant we obtain

$$X_t = X_0 + \int_0^t K(t-s)(b^0 + BX_s)ds + \int_0^t K(t-s)\sigma dW_s$$

Volterra–Ornstein–Uhlenbeck process

- ▶ With $E = \mathbb{R}^d$ and $\sigma(x) \equiv \sigma$ constant we obtain

$$X_t = X_0 + \int_0^t K(t-s)(b^0 + BX_s)ds + \int_0^t K(t-s)\sigma dW_s$$

- ▶ This is a **Gaussian process**.

Volterra–Ornstein–Uhlenbeck process

- ▶ With $E = \mathbb{R}^d$ and $\sigma(x) \equiv \sigma$ constant we obtain

$$X_t = X_0 + \int_0^t K(t-s)(b^0 + BX_s)ds + \int_0^t K(t-s)\sigma dW_s$$

- ▶ This is a **Gaussian process**.
- ▶ The Riccati–Volterra equation has an **explicit solution**:

$$\psi = uE_B + f * E_B.$$

where R_B is the resolvent of $-KB$ and $E_B = K - R_B * K$.

Volterra–Ornstein–Uhlenbeck process

- ▶ With $E = \mathbb{R}^d$ and $\sigma(x) \equiv \sigma$ constant we obtain

$$X_t = X_0 + \int_0^t K(t-s)(b^0 + BX_s)ds + \int_0^t K(t-s)\sigma dW_s$$

- ▶ This is a **Gaussian process**.
- ▶ The Riccati–Volterra equation has an **explicit solution**:

$$\psi = uE_B + f * E_B.$$

where R_B is the resolvent of $-KB$ and $E_B = K - R_B * K$.

- ▶ The quadratic variation of the process Y is deterministic,

$$\langle Y \rangle_t = \int_0^t \psi(T-s)\sigma\sigma^\top\psi(T-s)^\top ds.$$

Thus the martingale condition holds.

The martingale condition

How to verify the martingale property of e^{Y_t} more generally?

- ▶ In the **classical case** ($K \equiv \text{id}$) the condition is

$$\operatorname{Re} \phi(t) + \operatorname{Re} \psi(t)x \leq 0, \quad t \geq 0, \quad x \in E.$$

- ▶ Then

$$\operatorname{Re} Y_t = \operatorname{Re} \phi(t) + \operatorname{Re} \psi(t)X_t \leq 0,$$

so that e^{Y_t} is bounded.

The martingale condition

- ▶ A **resolvent of the first kind** of K is a kernel L such that

$$K * L = L * K \equiv \text{id}$$

In general L is a measure, for instance $L(dt) = \delta_0(dt)$ if $K \equiv \text{id}$.

The martingale condition

- ▶ A **resolvent of the first kind** of K is a kernel L such that

$$K * L = L * K \equiv \text{id}$$

In general L is a measure, for instance $L(dt) = \delta_0(dt)$ if $K \equiv \text{id}$.

- ▶ **Example:** If $K(t) = \frac{1}{\Gamma(\alpha)} t^{\alpha-1}$, $\alpha \in (\frac{1}{2}, 1)$, then

$$L(t) = \frac{1}{\Gamma(1-\alpha)} t^{-\alpha}$$

The martingale condition

- ▶ A **resolvent of the first kind** of K is a kernel L such that

$$K * L = L * K \equiv \text{id}$$

In general L is a measure, for instance $L(dt) = \delta_0(dt)$ if $K \equiv \text{id}$.

- ▶ **Example:** If $K(t) = \frac{1}{\Gamma(\alpha)} t^{\alpha-1}$, $\alpha \in (\frac{1}{2}, 1)$, then

$$L(t) = \frac{1}{\Gamma(1-\alpha)} t^{-\alpha}$$

- ▶ **Example:** If K is **completely monotone**, then L exists and is the sum of a point mass in zero and a completely monotone function.

The martingale condition

Shift operator: $\Delta_\tau f(t) = f(t + \tau)$

The martingale condition

Shift operator: $\Delta_\tau f(t) = f(t + \tau)$

Lemma. Consider the setting of Theorem (*), and assume K admits a resolvent of the first kind L . Define

$$\pi_\tau = \Delta_\tau \psi * L - \Delta_\tau(\psi * L),$$

and assume $\pi_\tau \in BV_{\text{loc}}$ for every $\tau \geq 0$. Then

$$Y_t = \phi(\tau) + (\Delta_\tau \psi * L)(0)X_t - \pi_\tau(t)X_0 + (d\pi_\tau * X)_t,$$

with $\tau = T - t$ and

$$\phi(\tau) = \int_0^\tau \left(\psi(s)b^0 + \frac{1}{2}\psi(s)A^0\psi(s)^\top \right) ds.$$

The martingale condition

Shift operator: $\Delta_\tau f(t) = f(t + \tau)$

Lemma. Consider the setting of Theorem (*), and assume K admits a resolvent of the first kind L . Define

$$\pi_\tau = \Delta_\tau \psi * L - \Delta_\tau (\psi * L),$$

and assume $\pi_\tau \in BV_{\text{loc}}$ for every $\tau \geq 0$. Then

$$Y_t = \phi(\tau) + (\Delta_\tau \psi * L)(0)X_t - \pi_\tau(t)X_0 + (d\pi_\tau * X)_t,$$

with $\tau = T - t$ and

$$\phi(\tau) = \int_0^\tau \left(\psi(s)b^0 + \frac{1}{2}\psi(s)A^0\psi(s)^\top \right) ds.$$

In particular: For $E = \mathbb{R}_+^d$, verify martingale condition by controlling the signs of the real parts of $(\Delta_\tau \psi * L)(0)$, $\pi_\tau(t)$, and $d\pi_\tau$.

Volterra square-root process

- ▶ $E = \mathbb{R}_+^d$ and diagonal kernel $K = \text{diag}(K_1, \dots, K_d)$:

$$X_{i,t} = X_{i,0} + \int_0^t K_i(t-s)b_i(X_s)ds + \int_0^t K_i(t-s)\sigma_i\sqrt{X_{i,s}}dW_{i,s}$$

- ▶ Inward-pointing drift condition:

$$b^0 \in \mathbb{R}_+^d \text{ and } B_{ij} \geq 0 \text{ for } i \neq j.$$

Volterra square-root process

- ▶ $E = \mathbb{R}_+^d$ and diagonal kernel $K = \text{diag}(K_1, \dots, K_d)$:

$$X_{i,t} = X_{i,0} + \int_0^t K_i(t-s)b_i(X_s)ds + \int_0^t K_i(t-s)\sigma_i\sqrt{X_{i,s}}dW_{i,s}$$

- ▶ Inward-pointing drift condition:

$$b^0 \in \mathbb{R}_+^d \text{ and } B_{ij} \geq 0 \text{ for } i \neq j.$$

- ▶ **Assumption:** Each K_i is **completely monotone** and is controlled near zero in the sense that

$$K_i(t) = o(t^{-\gamma_i}) \text{ as } t \rightarrow 0$$

for some $\gamma_i < 1/2$ (these assumptions can be relaxed).

The multi-factor Volterra CIR process

Theorem.

- ▶ The stochastic Volterra equation has a unique in law \mathbb{R}_+^d -valued weak solution for any initial condition $X_0 \in \mathbb{R}_+^d$. The paths of X_i are Hölder continuous of any order less than $H_i = 1/2 - \gamma_i$, for each $i = 1, \dots, d$.
- ▶ For any $u \in (\mathbb{C}^d)^*$ with $\operatorname{Re} u_i \leq 0$ for each $i = 1, \dots, d$, the Riccati–Volterra equation

$$\psi_i(t) = u_i K_i(t) + \int_0^t K_i(t-s) \left(\psi(s) b^i + \frac{\sigma_i^2}{2} \psi_i(s)^2 \right) ds$$

has a unique global solution $\psi \in L_{\text{loc}}^2(\mathbb{R}_+, (\mathbb{C}^d)^*)$.

- ▶ The martingale condition in Theorem (*) holds, as does the affine transform formula.

Conclusion

- ▶ Brownian paths are too smooth for volatility modeling
- ▶ Affine Volterra processes generalize various known rough volatility models
- ▶ Despite lack of Markov property, affine transform formulas can be derived
- ▶ Lots to do:
 - ▶ Numerical methods for the Riccati–Volterra equations . . .
 - ▶ . . . or even explicit solutions in special cases?
 - ▶ Hedging and optimal investment in these models
 - ▶ Boundary attainment for Volterra square-root processes
 - ▶ Non-convolution kernels $K(t, s)$
 - ▶ Etc.

Thank you!