

Maximum likelihood estimation for random walks in random environments

F. Comets, M. Falconnet, A. Gloter, O. Loukianov, D. Loukianova, C. Mattias

SAPS 2017, Peterhof

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties
- Construction

Addendum

- Numerical simulations
- Extensions

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties
- Construction

Addendum

- Numerical simulations
- Extensions

Framework: We consider a one-dimensional random walk in i.i.d. random environment (RWRE) with a parametric distribution.

Result: Based on a single observation of the path, we provide a maximum likelihood estimation procedure for the law of the environment.

Random environment on \mathbb{Z}

- ▶ $\omega = \{\omega_x\}_{x \in \mathbb{Z}}$ i.i.d. with $\omega_x \in]0, 1[$ and $\omega_x \sim \mu$,
- ▶ $\mathbb{P} = \mu^{\otimes \mathbb{Z}}$ law on $]0, 1[^{\mathbb{Z}}$ of ω and \mathbb{E} expectation

Markov process conditional on the environment

For *fixed* ω , let $X = \{X_t\}_{t \in \mathbb{N}}$ be the Markov chain on \mathbb{Z} starting at $X_0 = 0$ and with transitions

$$P_\omega(X_{t+1} = y | X_t = x) = \begin{cases} \omega_x & \text{if } y = x + 1 \\ 1 - \omega_x & \text{if } y = x - 1 \end{cases}$$

P_ω is the measure on the path space of X given ω (*quenched* law) and E_ω corresponding expectation.

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Random environment on \mathbb{Z}

- ▶ $\omega = \{\omega_x\}_{x \in \mathbb{Z}}$ i.i.d. with $\omega_x \in]0, 1[$ and $\omega_x \sim \mu$,
- ▶ $\mathbb{P} = \mu^{\otimes \mathbb{Z}}$ law on $]0, 1[^{\mathbb{Z}}$ of ω and \mathbb{E} expectation

Markov process conditional on the environment

For *fixed* ω , let $X = \{X_t\}_{t \in \mathbb{N}}$ be the Markov chain on \mathbb{Z} starting at $X_0 = 0$ and with transitions

$$P_\omega(X_{t+1} = y | X_t = x) = \begin{cases} \omega_x & \text{if } y = x + 1 \\ 1 - \omega_x & \text{if } y = x - 1 \end{cases}$$

P_ω is the measure on the path space of X given ω (*quenched law*) and E_ω corresponding expectation.

[Introduction](#)[Framework](#)[Problem](#)[MLE construction](#)[Transient case](#)[Properties](#)[Branching process](#)[Results](#)[Recurrent case](#)[Assumptions](#)[Properties](#)[Construction](#)[Addendum](#)[Numerical simulations](#)[Extensions](#)

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Random walk in random environment (RWRE)

The (unconditional) law of X is the *annealed* law

$$\mathbf{P}(\cdot) = \mathbb{E}(P_{\omega}(\cdot)) = \int P_{\omega}(\cdot) d\mathbb{P}(\omega),$$

with \mathbf{E} the corresponding expectation.

Note that X is *not* a Markov process under \mathbf{P} in general.

Consider the "left/right" ratio

$$\rho_x = \frac{1 - \omega_x}{\omega_x}, \quad x \in \mathbb{Z}$$

Solomon(1975) has proved the classification:

Recurrent case

If $\mathbb{E}(\log \rho_0) = 0$, then

$$-\infty = \liminf_{t \rightarrow \infty} X_t < \limsup_{t \rightarrow \infty} X_t = +\infty, \quad \mathbf{P}\text{-a.s.}$$

and X_t is null-recurrent.

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Transient case

if $\mathbb{E}(\log \rho_0) < 0$, then

$$\lim_{t \rightarrow \infty} X_t = +\infty, \quad \mathbf{P}\text{-a.s.}$$

Moreover, if $T_n = \inf\{t \in \mathbb{N} : X_t = n\}$, then

- ▶ **Ballistic case:** if $\mathbb{E}(\rho_0) < 1$, then $T_n/n \rightarrow c$ \mathbf{P} -a.s. when $n \rightarrow \infty$.
- ▶ **Sub-ballistic case:** If $\mathbb{E}(\rho_0) \geq 1$ and $\mathbb{E}(\rho_0^\kappa) = 1$ for some $0 < \kappa \leq 1$ then (in general) $T_n \sim n^{1/\kappa}$.

Environment law estimation

Estimate μ from a single observation $(X_t)_{0 \leq t \leq T}$ of a RWRE path.

Assumptions

We suppose that $\mu = \mu_{\theta^*} \in \{\mu_{\theta}\}_{\theta \in \Theta}$, where $\theta^* \in \Theta$ is an unknown parameter, $\Theta \subset \mathbb{R}^d$ compact.

Example (finitely supported law)

$$\mu(\{a_i\}, \{p_i\}) = \sum_{i=1}^m p_i \delta_{a_i}, \quad \mathbb{E} \rho_0 = \sum_{i=1}^m p_i \log \frac{1 - a_i}{a_i}.$$

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Environment law estimation

Estimate μ from a single observation $(X_t)_{0 \leq t \leq T}$ of a RWRE path.

Assumptions

We suppose that $\mu = \mu_{\theta^*} \in \{\mu_{\theta}\}_{\theta \in \Theta}$, where $\theta^* \in \Theta$ is an unknown parameter, $\Theta \subset \mathbb{R}^d$ compact.

Example (finitely supported law)

$$\mu(\{a_i\}, \{p_i\}) = \sum_{i=1}^m p_i \delta_{a_i}, \quad \mathbb{E} \rho_0 = \sum_{i=1}^m p_i \log \frac{1 - a_i}{a_i}.$$

We write \mathbb{P}^{θ} , \mathbf{P}^{θ} and so on for RWRE generated by μ_{θ} , and \mathbb{P}^{\star} , \mathbf{P}^{\star} , ... for $\theta = \theta^*$ (the true parameter to estimate).

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Environment law estimation

Estimate μ from a single observation $(X_t)_{0 \leq t \leq T}$ of a RWRE path.

Assumptions

We suppose that $\mu = \mu_{\theta^*} \in \{\mu_{\theta}\}_{\theta \in \Theta}$, where $\theta^* \in \Theta$ is an unknown parameter, $\Theta \subset \mathbb{R}^d$ compact.

Example (finitely supported law)

$$\mu(\{a_i\}, \{p_i\}) = \sum_{i=1}^m p_i \delta_{a_i}, \quad \mathbb{E} \rho_0 = \sum_{i=1}^m p_i \log \frac{1 - a_i}{a_i}.$$

We write \mathbb{P}^{θ} , \mathbf{P}^{θ} and so on for RWRE generated by μ_{θ} , and \mathbb{P}^* , \mathbf{P}^* , ... for $\theta = \theta^*$ (the true parameter to estimate).

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

O. Adelman and N. Enriquez (2004), Random walks in random environment: what a single trajectory tells.

A nice family of estimators of *moments* of μ_{θ^*} .

Example: first steps from each site \Rightarrow first moment.

Drawback:

- ▶ Which moments to estimate to recover μ_{θ^*} ?
- ▶ Only some steps are used (loss of information).

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Maximum likelihood estimator

Fix a time T , a trajectory $X_{[0,T]}$, and let $L_x = L_x(T)$ and $R_x = R_x(T)$ be the number of left and right steps from site x . Then,

$$P_\omega(X_{[0,T]}) = \prod_{x \in \mathbb{Z}} \omega_x^{R_x} (1 - \omega_x)^{L_x}$$

and

$$\mathbf{P}^\theta(X_{[0,T]}) = \mathbb{E}^\theta \prod_{x \in \mathbb{Z}} \omega_x^{R_x} (1 - \omega_x)^{L_x} = \prod_{x \in \mathbb{Z}} \mathbb{E}^\theta \omega_x^{R_x} (1 - \omega_x)^{L_x} = \prod_{x \in \mathbb{Z}} \int_0^1 a^{R_x} (1 - a)^{L_x} d\mu_\theta(a).$$

Introduction

[Framework](#)[Problem](#)[MLE construction](#)

Transient case

[Properties](#)[Branching process](#)[Results](#)

Recurrent case

[Assumptions](#)[Properties](#)[Construction](#)

Addendum

[Numerical simulations](#)[Extensions](#)

Let ϕ_θ be the function from \mathbb{N}^2 to \mathbb{R} given by

$$\phi_\theta(x, y) = \log \int_0^1 a^x (1-a)^y d\mu_\theta(a).$$

The criterion function $\theta \mapsto \ell_T(\theta)$ is defined as

$$\ell_T(\theta) = \log \mathbf{P}^\theta(X_{[0, T]}) = \sum_{x \in \mathbb{Z}} \phi_\theta(R_x, L_x),$$

and our estimator is

$$\hat{\theta}_T \in \operatorname{Argmax}_{\theta \in \Theta} \ell_T(\theta).$$

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Study the convergence of $\hat{\theta}_T$ to θ^* .

Method: Show that $\ell_T(\theta)$ converges (after appropriate normalisation) to some $\ell(\theta)$ with

$$\operatorname{Argmax}_{\theta \in \Theta} \ell(\theta) = \theta^*$$

and apply classical M -estimation theory.

Question: where $\ell(\theta)$ comes from?

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Introduction

Framework
Problem
MLE construction

Transient case

Properties
Branching process
Results

Recurrent case

Assumptions
Properties
Construction

Addendum

Numerical simulations
Extensions

We take $n \in \mathbb{N}$ and $T = T_n = \inf\{t \in \mathbb{N} : X_t = n\}$.

Note that

- ▶ Only the visited sites contribute to $\ell_{T_n}(\theta)$.
- ▶ The number of visited sites $x < 0$ is bounded (since X is transient to the right).
- ▶ Moreover, $R_x = L_{x+1} + 1$ for $x = 0, 1, \dots, n-1$.

Hence

$$\ell_{T_n}(\theta) \approx \sum_{x=0}^{n-1} \phi_{\theta}(L_{x+1} + 1, L_x).$$

Under \mathbf{P}^θ , the sequence L_n, L_{n-1}, \dots, L_0 has the same distribution as a BPI denoted Z_0, \dots, Z_n , and defined by

$$Z_0 = 0 \quad \text{and} \quad Z_{k+1} = \sum_{i=0}^{Z_k} \xi_{k+1,i} \quad \text{for } k \geq 0,$$

with $\{\xi_{k,i}\}$ independent and

$$\forall m \in \mathbb{N}, \quad P_\omega(\xi_{k,i} = m) = (1 - \omega_k)^m \omega_k,$$

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Under \mathbf{P}^θ , $\{Z_n\}$ is an irreducible positive recurrent homogeneous Markov chain with the transition kernel

$$Q_\theta(x, y) = \binom{x+y}{x} \int_0^1 a^{x+1} (1-a)^y d\mu_\theta(a).$$

Consequence

$$\frac{1}{n} \ell_{T_n}(\theta) \sim \frac{1}{n} \sum_{k=0}^{n-1} \phi_\theta(Z_k + 1, Z_{k+1}) \quad \text{under } \mathbf{P}^\star$$

and the right-hand side is (up to constants) the likelihood of a Markov process.

It follows that $\phi_\theta(Z_k + 1, Z_{k+1})$ satisfies a law of large numbers.

Introduction

[Framework](#)[Problem](#)[MLE construction](#)

Transient case

[Properties](#)[Branching process](#)[Results](#)

Recurrent case

[Assumptions](#)[Properties](#)[Construction](#)

Addendum

[Numerical simulations](#)[Extensions](#)

We deduce that $\ell_{T_n}(\theta)/n$ converges in \mathbf{P}^* -probability to a deterministic limit $\ell(\theta)$:

$$\ell(\theta) = \mathbf{E}^* \phi_\theta(\tilde{Z}_0 + 1, \tilde{Z}_1)$$

Ballistic case

\tilde{Z}_k has a finite first order moment.

Sub-ballistic case

We fix $\theta_0 \in \Theta$ and replace $\ell_{T_n}(\theta)$ with

$$\ell_{T_n}(\theta) - \ell_{T_n}(\theta_0) \sim \sum_{k=0}^{n-1} (\phi_\theta(Z_k + 1, Z_{k+1}) - \phi_{\theta_0}(Z_k + 1, Z_{k+1}))$$

and assume that $\phi_\theta - \phi_{\theta_0}$ is uniformly integrable (true in most cases).

Using the almost linear nature of ϕ_θ , we prove that $\ell(\theta)$ is finite, with a maximum at θ^* .

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Results: consistency, asymptotic normality and efficiency

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

The standard M -estimators theory then applies. Under appropriate (classical) assumptions, in the *transient* case, we establish that MLE satisfies

- ▶ **Consistency:** $\lim_{n \rightarrow +\infty} \hat{\theta}_{T_n} = \theta^*$, \mathbf{P}^* -a.s.
- ▶ **Asymptotic normality:**
 $\sqrt{n}(\hat{\theta}_{T_n} - \theta^*) \rightsquigarrow^{\mathbf{P}^* - dist.} \mathcal{N}(0, \Sigma_{\theta^*}^{-1}).$
- ▶ **Efficiency:** Σ_{θ^*} is the Fisher information matrix.

Hence the rate of convergence is of the order \sqrt{T} in the ballistic case, and $T^{\kappa/2}$ in the sub-ballistic case ($\kappa \leq 1$).

Recurrent case

We consider the distributions of the form

$$\mu_{(\mathbf{a}, \mathbf{p})} = \sum_{i=1}^d p_i \delta_{a_i}$$

and assume that the true parameter $\theta^* = (\mathbf{a}^*, \mathbf{p}^*)$ belongs to a compact $\Theta \subset (0, 1)^{2d}$ satisfying

Assumption (Identifiability)

For any $\theta = (\mathbf{a}, \mathbf{p})$ in Θ ,

$$0 < a_1 < a_2 < \dots < a_d < 1$$

Assumption (Recurrent environment)

$$\mathbb{E}^* \rho_0 = \sum_{i=1}^d p_i^* \log \frac{1 - a_i^*}{a_i^*} = 0.$$

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions


Example (Temkin)

Let


$$\mu_\theta = \frac{1}{2}\delta_a + \frac{1}{2}\delta_{1-a}.$$

Here, the unknown parameter is $a \in \Theta \subset (0, 1/2)$.

Energy landscape



Trajectory



Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Properties of recurrent RWRE

An important property of recurrent RWRE is *localization*: the RE creates *traps* where the walk spends much time.

A useful trap visualization is the potential landscape V where $V = \{V(x) : x \in \mathbb{Z}\}$ is defined by

$$V(x) = \begin{cases} \sum_{y=0}^x \log \rho_y - \log \rho_0 & \text{if } x \geq 0 \\ -\sum_{y=x+1}^0 \log \rho_y & \text{if } x < 0 \end{cases}$$

The environment $\{\omega_x\}$ can be recovered from its potential:

$$\omega_x = \frac{\exp(-V(x))}{\exp(-V(x)) + \exp(-V(x-1))}.$$

Introduction

[Framework](#)[Problem](#)[MLE construction](#)

Transient case

[Properties](#)[Branching process](#)[Results](#)

Recurrent case

[Assumptions](#)**[Properties](#)**[Construction](#)

Addendum

[Numerical simulations](#)[Extensions](#)

Main valleys




Figure :

$$c_n = \min \{x \geq 0 : V(x) - \min_{0 \leq y \leq x} V(y) \geq \log n + (\log n)^{1/2}\},$$

$$b_n = \min \{x \geq 0 : V(x) = \min_{0 < y < c_n} V(y)\}.$$

Basic localization properties of the RW are known since the works of Sinai (1982), Golosov (1984) and others. Namely,

- ▶ with an overwhelming probability, the (reflected) walk $X_{[0,n]}$ stays between 0 and c_n (Arrhenius law);
- ▶ $b_n/\log^2 n$ and $c_n/\log^2 n$ converge in law to some non-degenerate random variables;
- ▶ moreover, $(X_n - b_n)/\log^2 n$ converges to 0 in probability.

For MLE, we need to describe the distribution of local times of $X_n - b_n$.

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Let $\tilde{V} = \{\tilde{V}(x) : x \in \mathbb{Z}\}$ be a collection of random variables distributed as V “conditioned” to stay positive. For each \tilde{V} , let $\tilde{\omega}$ be the corresponding environment on \mathbb{Z} .

Let $\nu(x) = \nu^+(x) + \nu^-(x)$ be the invariant measure of the corresponding (ergodic) Markov chain \tilde{X}_n on \mathbb{Z} , where

$$\nu^+(x) = \frac{e^{-\tilde{V}(x)}}{2 \sum_{z \in \mathbb{Z}} e^{-\tilde{V}(z)}} \quad \text{and} \quad \nu^-(x) = \frac{e^{-\tilde{V}(x-1)}}{2 \sum_{z \in \mathbb{Z}} e^{-\tilde{V}(z)}}.$$

Then $\tilde{\omega}_x = \nu^+(x)/\nu(x)$.

Remark

The possible values of $\tilde{\omega}$ are those of ω , though their distributions are different (not i.i.d.)

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties**
- Construction

Addendum

- Numerical simulations
- Extensions

Gantert-Peres-Shi theorem

Put $\nu_n^+(x) = R_x/n$, $\nu_n^-(x) = L_x/n$.

Theorem (Gantert-Peres-Shi, 2010)

The distributions of

$$\{(\nu_n^+(x + b_n), \nu_n^-(x + b_n)) : x \in \mathbb{Z}\}$$

converge weakly to the distribution of

$$\{(\nu^+(x), \nu^-(x)) : x \in \mathbb{Z}\}.$$

As a consequence, for each strongly continuous functional f which is translation invariant, we have

$$f(\{(\nu_n^+, \nu_n^-)\}) \xrightarrow[n \rightarrow \infty]{law} f(\{(\nu^+, \nu^-)\}).$$

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

The annealed log-likelihood $\ell_n(\theta) = \log \mathbf{P}^\theta(X_{[0,n]})$ in our case is given by

$$\ell_n(\theta) = \sum_{x \in \mathbb{Z}} \log \left[\sum_{i=1}^d a_i^{R_x} (1 - a_i)^{L_x} p_i \right]$$

Recurrence imply $R_x, L_x \rightarrow \infty$ as $n \rightarrow \infty$, so

- ▶ the branching explodes;
- ▶ but we can apply Laplace methods.

Denote by \mathcal{R}_n the range of the walk:

$$\mathcal{R}_n = \{x : \exists t \leq n, X_t = x\}$$

Recall that $|\mathcal{R}_n| = O_{\mathbb{P}^*}(\log^2 n)$.

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Log-likelihood decomposition

For any x in \mathcal{R}_n , define the random integer

$$\hat{i} = \hat{i}(\mathbf{a}, n, x) = \operatorname{Argmax}_i \left\{ a_i^{R_x} (1 - a_i)^{L_x} \right\}$$

Then

$$\begin{aligned} \ell_n(\theta) &= \sum_{x \in \mathbb{Z}} (R_x \log a_{\hat{i}} + L_x \log(1 - a_{\hat{i}})) + \sum_{x \in \mathcal{R}_n} \log p_{\hat{i}} \\ &\quad + \sum_{x \in \mathcal{R}_n} \log \left(1 + \sum_{i \neq \hat{i}} \left(\frac{a_i}{a_{\hat{i}}} \right)^{R_x} \left(\frac{1 - a_i}{1 - a_{\hat{i}}} \right)^{L_x} \frac{p_i}{p_{\hat{i}}} \right) \\ &= M_n + K_n + r_n = O(n) + O_{\mathbb{P}^*}(\log^2 n) + o_{\mathbb{P}^*}(\log^2 n) \end{aligned}$$

Introduction

[Framework](#)[Problem](#)[MLE construction](#)

Transient case

[Properties](#)[Branching process](#)[Results](#)

Recurrent case

[Assumptions](#)[Properties](#)[Construction](#)

Addendum

[Numerical simulations](#)[Extensions](#)

We define a MLE as

$$\hat{\theta}_n = (\hat{\mathbf{a}}_n, \hat{\mathbf{p}}_n) = \underset{(\mathbf{a}, \mathbf{p}) \in \Theta}{\operatorname{Argmax}} \ell_n(\theta),$$

and a (pseudo) MLE $(\overline{\mathbf{a}}_n, \overline{\mathbf{p}}_n)$ as

$$\begin{cases} \overline{\mathbf{a}}_n &= \underset{\mathbf{a} \in \Theta_a}{\operatorname{Argmax}} M_n(\mathbf{a}), \\ \overline{\mathbf{p}}_n &= \underset{\mathbf{p}}{\operatorname{Argmax}} K_n(\overline{\mathbf{a}}_n, \mathbf{p}). \end{cases}$$

Theorem

Both the the MPL estimator $(\overline{\mathbf{a}}_n, \overline{\mathbf{p}}_n)$ and ML estimator $(\hat{\mathbf{a}}_n, \hat{\mathbf{p}}_n)$ converge in \mathbf{P}^ -probability to the true parameter value θ^* .*

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties
- Construction

Addendum

- Numerical simulations
- Extensions

We define a MLE as

$$\hat{\theta}_n = (\hat{\mathbf{a}}_n, \hat{\mathbf{p}}_n) = \underset{(\mathbf{a}, \mathbf{p}) \in \Theta}{\operatorname{Argmax}} \ell_n(\theta),$$

and a (pseudo) MLE $(\overline{\mathbf{a}}_n, \overline{\mathbf{p}}_n)$ as

$$\begin{cases} \overline{\mathbf{a}}_n &= \underset{\mathbf{a} \in \Theta_a}{\operatorname{Argmax}} M_n(\mathbf{a}), \\ \overline{\mathbf{p}}_n &= \underset{\mathbf{p}}{\operatorname{Argmax}} K_n(\overline{\mathbf{a}}_n, \mathbf{p}). \end{cases}$$

Theorem

Both the the MPL estimator $(\overline{\mathbf{a}}_n, \overline{\mathbf{p}}_n)$ and ML estimator $(\hat{\mathbf{a}}_n, \hat{\mathbf{p}}_n)$ converge in \mathbf{P}^ -probability to the true parameter value θ^* .*

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Recall the properties of the *cross-entropy* $H(\mathbf{p}, \mathbf{q})$ of two (finitely supported) probability measures:

$$H(\mathbf{p}, \mathbf{q}) = -E_{\mathbf{p}} \log \mathbf{q} = - \sum_i p_i \log q_i$$

$$H(\mathbf{p}) = H(\mathbf{p}, \mathbf{p}) < H(\mathbf{p}, \mathbf{q}) \quad \text{if} \quad \mathbf{p} \neq \mathbf{q}$$

In particular, for $0 < p, q < 1$

$$\begin{aligned} \max_q \{p \log q + (1-p) \log(1-q)\} \\ = p \log p + (1-p) \log(1-p) \end{aligned}$$

Introduction

Framework

Problem

MLE construction

Transient case

Properties

Branching process

Results

Recurrent case

Assumptions

Properties

Construction

Addendum

Numerical simulations

Extensions

Recall that $\nu_n^+(x) = R_x/n$, $\nu_n^- = L_x/n$ and $\tilde{\omega} = \nu^+/\nu$. So

$$M_n = n \sum_{x \in \mathbb{Z}} \max_i \{ \nu_n^+(x) \log a_i + \nu_n^-(x) \log(1 - a_i) \}$$

and GPS theorem yields

$$\begin{aligned} \frac{M_n}{n} &\xrightarrow[n \rightarrow \infty]{\text{law}} M(\mathbf{a}, \nu^+, \nu^-) \\ &= \sum_{x \in \mathbb{Z}} \nu(x) \max_i \{ \tilde{\omega}_x \log a_i + (1 - \tilde{\omega}_x) \log(1 - a_i) \} \\ &= - \sum_{x \in \mathbb{Z}} \nu(x) \max_i H(\tilde{\omega}_x, a_i) \end{aligned}$$

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties
- Construction

Addendum

- Numerical simulations
- Extensions

Using $\tilde{\omega}_x \in \{a_i^*\}$, it's easily seen that for $\mathbf{a} \neq \mathbf{a}^*$,

$$M(\mathbf{a}) < M(\mathbf{a}^*) = - \sum_{x \in \mathbb{Z}} \nu(x) H(\tilde{\omega}_x).$$

Finally,

$$\frac{M_n(\mathbf{a}^*) - M_n(\mathbf{a})}{n} \xrightarrow[n \rightarrow \infty]{\text{law}} M(\mathbf{a}^*) - M(\mathbf{a}) > 0$$

whence it can be deduced that

$$\operatorname{Argmax}_{\mathbf{a} \in \Theta} M_n(\mathbf{a}) = \overline{\mathbf{a}_n} \rightarrow \mathbf{a}^*.$$

Introduction

Framework
Problem
MLE construction

Transient case

Properties
Branching process
Results

Recurrent case

Assumptions
Properties
Construction

Addendum

Numerical simulations
Extensions

Probability estimation

Recall the definition of $\overline{\mathbf{p}}_n$:

$$\overline{\mathbf{p}}_n = \underset{\mathbf{p}}{\operatorname{Argmax}} \sum_{x \in \mathcal{R}_n} \log p_{\hat{i}} = \underset{\mathbf{p}}{\operatorname{Argmax}} \sum_{i=1}^d \frac{|\mathcal{R}_n(i)|}{|\mathcal{R}_n|} \log p_i,$$

where

$$\hat{i} = \underset{i}{\operatorname{Argmax}} \{R_x \log(\overline{\mathbf{a}}_n)_i + L_x \log(1 - (\overline{\mathbf{a}}_n)_i)\}$$

and $\mathcal{R}_n(i) = \{x \in \mathcal{R}_n : \hat{i} = i\}$.

By the law of large numbers, $R_x/(R_x + L_x) \rightarrow \omega_x$, hence $\omega_x = a_{\hat{i}}^*$ for $\mathbf{a} \approx \mathbf{a}^*$ if n is large enough.

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties
- Construction

Addendum

- Numerical simulations
- Extensions

Probability estimation

Recall the definition of $\overline{\mathbf{p}}_n$:

$$\overline{\mathbf{p}}_n = \underset{\mathbf{p}}{\operatorname{Argmax}} \sum_{x \in \mathcal{R}_n} \log p_{\hat{i}} = \underset{\mathbf{p}}{\operatorname{Argmax}} \sum_{i=1}^d \frac{|\mathcal{R}_n(i)|}{|\mathcal{R}_n|} \log p_i,$$

where

$$\hat{i} = \underset{i}{\operatorname{Argmax}} \{R_x \log(\overline{\mathbf{a}}_n)_i + L_x \log(1 - (\overline{\mathbf{a}}_n)_i)\}$$

and $\mathcal{R}_n(i) = \{x \in \mathcal{R}_n : \hat{i} = i\}$.

By the law of large numbers, $R_x/(R_x + L_x) \rightarrow \omega_x$, hence $\omega_x = a_i^*$ for $\mathbf{a} \approx \mathbf{a}^*$ if n is large enough.

Since $\overline{\mathbf{a}}_n \rightarrow \mathbf{a}^*$, we get $|\mathcal{R}_n(i)| \rightarrow \#\{x \in \mathcal{R}_n : \omega_x = a_i^*\}$, whence $|\mathcal{R}_n(\cdot)|/|\mathcal{R}_n| = \overline{\mathbf{p}}_n \rightarrow \mathbf{p}^*$.

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties
- Construction

Addendum

- Numerical simulations
- Extensions

Probability estimation

Recall the definition of $\overline{\mathbf{p}}_n$:

$$\overline{\mathbf{p}}_n = \underset{\mathbf{p}}{\operatorname{Argmax}} \sum_{x \in \mathcal{R}_n} \log p_{\hat{i}} = \underset{\mathbf{p}}{\operatorname{Argmax}} \sum_{i=1}^d \frac{|\mathcal{R}_n(i)|}{|\mathcal{R}_n|} \log p_i,$$

where

$$\hat{i} = \underset{i}{\operatorname{Argmax}} \{R_x \log(\overline{\mathbf{a}}_n)_i + L_x \log(1 - (\overline{\mathbf{a}}_n)_i)\}$$

and $\mathcal{R}_n(i) = \{x \in \mathcal{R}_n : \hat{i} = i\}$.

By the law of large numbers, $R_x/(R_x + L_x) \rightarrow \omega_x$, hence $\omega_x = a_{\hat{i}}^*$ for $\mathbf{a} \approx \mathbf{a}^*$ if n is large enough.

Since $\overline{\mathbf{a}}_n \rightarrow \mathbf{a}^*$, we get $|\mathcal{R}_n(i)| \rightarrow \#\{x \in \mathcal{R}_n : \omega_x = a_i^*\}$, whence $|\mathcal{R}_n(\cdot)|/|\mathcal{R}_n| = \overline{\mathbf{p}}_n \rightarrow \mathbf{p}^*$.

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties
- Construction

Addendum

- Numerical simulations
- Extensions

Example (Temkin)

Introduction

- Framework
- Problem
- MLE construction

Transient case

- Properties
- Branching process
- Results

Recurrent case

- Assumptions
- Properties
- Construction

Addendum

- Numerical simulations
- Extensions




Figure : Boxplots of our estimator (white) and Adelman and Enriquez estimator (grey). The true value of θ^* is 0.3.

Introduction

Framework
Problem
MLE construction

Transient case

Properties
Branching process
Results

Recurrent case

Assumptions
Properties
Construction

Addendum

Numerical simulations
Extensions

- ▶ P. Andreoletti, D. Loukianova, C. Matias (2015): Hidden Markov model for parameter estimation of a random walk in a Markov environment.
- ▶ R. Diel, M. Lerasle (2016): Non parametric estimation for random walks in random environment.